

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number  
**WO 01/04296 A1**

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
C07K 14/47, 16/18, A61K 38/17, 39/00

J4R 2G9 (CA). LA1, Albert [CA/CA]; 35 Wintermute  
Blvd., Scarborough, Ontario M1W 3M5 (CA).

(21) International Application Number: PCT/IB99/01690

(74) Agents: DUBUC, Jean, H. et al.; Goudreau Gage Dubuc  
& Martineau Walker, The Stock Exchange Tower, Suite  
3400, P.O. Box 242, 800 Place-Victoria, Montreal H4Z 1E9  
(CA).

(22) International Filing Date:  
24 September 1999 (24.09.1999)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/143,315 12 July 1999 (12.07.1999) US

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,  
BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE,  
ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP,  
KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,  
MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,  
SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ,  
VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier application:  
US 60/143,315 (CIP)  
Filed on 12 July 1999 (12.07.1999)

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): MCGILL  
UNIVERSITY [CA/CA]; 845 Sherbrooke Street West,  
Montreal, Quebec H3A 2T5 (CA).

Published:  
— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BRANTON, Philip,  
E. [CA/CA]; 266 Macaulay Street, St-Lambert, Quebec

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: RBP1 POLYPEPTIDES AND USES THEREOF

(57) Abstract: The invention features methods for modulating cell proliferation, methods for identifying compounds that modulate cell proliferation, and methods for determining whether a patient has a cell proliferation disease. The invention also features pharmaceutical compositions that include RBP1 polypeptides or nucleic acids, and uses thereof.

WO 01/04296 A1

Application No: 09/942,087  
Reference: E2

## RBP1 POLYPEPTIDES AND USES THEREOF

### Background of the Invention

5       The invention relates to the regulation of gene transcription and cell proliferation.

Control of gene expression by the E2F family of transcription factors plays an important role in the cell cycle and growth arrest, as E2F regulates expression of a number of genes required for DNA synthesis and cell cycle  
10   progression. E2F appears to be a major target of the retinoblastoma (RB) family of growth suppressors, pRB, p130 and p107, that bind to and inhibit E2F. Interactions between the E2F and RB families are cell cycle-dependent. pRB-E2F complexes are found throughout the cell cycle and in growth-arrested cells, and inhibition of E2F activity appears to be regulated at least in part by  
15   phosphorylation of pRB by cyclin-dependent protein kinases (Cdk). p107-E2F complexes are found largely in late G1 and S-phase, and also may regulate cell cycle progression. p130-E2F complexes are prominent in growth-arrested cells; in this context, the complexes have been shown to lack Cdk2 and cyclins.

### Summary of the Invention

20       In a first aspect, the invention features a method for modulating (e.g., decreasing) proliferation of a cell. The method includes contacting the cell with an RBP1 polypeptide that modulates proliferation of the cell. In one preferred embodiment, contacting is achieved by expressing in the cell a nucleic acid molecule encoding the RBP1 polypeptide.

25       In a second aspect, the invention features another method for modulating (e.g., decreasing) proliferation of a cell. The method includes contacting the cell with a compound that modulates RBP1 biological activity or expression.

In a third aspect, the invention features a method for identifying a

-2-

compound that modulates cell proliferation. The method generally includes contacting a cell with a candidate compound and monitoring the level of expression of an RBP1 polypeptide, wherein a change in the level of expression in response to the candidate compound, relative to a level of expression in a cell not contacted with the candidate compound, identifies the compound as a compound that modulates cell proliferation.

In a fourth aspect, the invention features another method for identifying a compound that modulates cell proliferation. The method includes: (a) providing an RBP1 polypeptide having biological activity, (b) contacting the polypeptide with a candidate compound, and (c) monitoring the level of the biological activity of the RBP1 polypeptide, wherein a change in the level of biological activity in response to the candidate compound, relative to a level of biological activity of an RBP1 polypeptide not contacted with the candidate compound, identifies the compound as a compound that modulates cell proliferation. The RBP1 polypeptide can be in a cell or in a cell-free system.

In a fifth aspect, the invention features a method for determining whether a first mammal (e.g., a human patient) has a cell proliferation disease or increased likelihood of developing the disease. The method includes measuring the level of expression of an RBP1 polypeptide in a sample from the mammal, wherein an increase in the level of expression of the RBP1 polypeptide, relative to a matched sample from a second mammal (e.g., an unaffected person) indicates that the first mammal has the cell proliferation disease or increased likelihood of developing the disease. In preferred embodiments, the expression is measured by assaying the amount of RBP1 polypeptide or RBP1 RNA in the sample.

In a sixth aspect, the invention features another method for determining whether a first mammal (e.g., a human patient) has a cell proliferation disease or an increased likelihood of developing the disease. The method includes

-3-

measuring the level of RBP1 biological activity in a sample from the mammal, wherein an alteration in the level of the biological activity, relative to a level of RBP1 biological activity in a matched sample from a second mammal (e.g., an unaffected person), indicates that the first mammal has the disease or increased  
5 likelihood of developing the disease.

In one preferred embodiment of the invention, the RBP1 biological activity is binding of the RBP1 polypeptide to an HDAC or to a protein (e.g., SAP30, SinA, or SinB) that is part of a protein complex containing an HDAC. Preferably, binding of the RBP1 polypeptide to the HDAC or the protein does  
10 not require the presence of an RB family member. In other preferred embodiments, the biological activity is binding of the RBP1 polypeptide to an E2F protein or is repression of E2F-mediated transcription. Preferably, the repression of E2F-mediated transcription does not require the presence of an HDAC protein. In particularly preferred embodiments of the invention, the cell  
15 is from a mammal (e.g., a human or a rodent) or in a mammal.

In a seventh aspect, the invention features a method of repressing transcription in a cell. The method includes introducing into the cell a polypeptide having: (i) a repressor domain from RBP1, and (ii) a DNA binding domain. In one preferred embodiment, this is achieved by expressing in the  
20 cell a nucleic acid molecule encoding the repressor domain from RBP1 and the DNA binding domain.

In an eighth aspect, the invention features a polypeptide that includes: (i) a repressor domain from RBP1, and (ii) a DNA binding domain. Preferably, the repressor domain includes amino acids 241-452 or amino acids 1167-1257  
25 of SEQ ID NO: 4.

In a ninth aspect, the invention features a nucleic acid molecule encoding a polypeptide having: (i) a repressor domain from RBP1, and (ii) a DNA binding domain.

-4-

In a tenth aspect, the invention features a composition that includes a substantially purified RBP1 polypeptide formulated in a physiologically acceptable carrier.

5 In an eleventh aspect, the invention features composition that includes a substantially purified antibody that specifically binds an RBP1 polypeptide formulated in a physiologically acceptable carrier.

10 In a twelfth aspect, the invention features a kit for diagnosing a mammal (e.g., a human) with a cell proliferation disease or an increased likelihood of developing the disease. The kit includes a substantially pure antibody that specifically binds an RBP1 polypeptide. Preferably, the kit also includes a means for detecting the binding of the antibody to the RBP1 polypeptide.

15 By "interact" or "bind" is meant that two polypeptides can transitorily bind, either directly or indirectly, to each other. Indirect binding can take place, for example, through the formation of a multi-protein complex. For example, HDACs are known to be part of a multi-protein complex that also includes, e.g., SAP30, SinA, and SinB (Pazin and Kadonaga, Cell 89:325-328, 1997). The interaction is dependent on inherent properties of the two polypeptides, and specifically excludes non-specific (e.g., strictly Brownian) contact. Preferably, this interaction can be disrupted or prevented by alteration in the primary, 20 secondary, or tertiary structure of one or both of the polypeptides.

By a "transcriptional repressor domain" is meant a polypeptide domain that retains its transcriptional repressor function when the domain is transferred to a heterologous protein. Preferably, the heterologous protein contains nuclear localization and DNA-binding domains.

25 By "RBP1 biological activity" is meant (i) RBP1 binding to an HDAC protein or a protein complex containing an HDAC, (ii) repression of E2F transcriptional activity, and (iii) arrest or decrease of cell cycle progression.

In addition to RBP1, other, related, polypeptides are highly likely to

-5-

have similar biological activities as RBP1, and, thus, are considered to be part of the invention. These polypeptides include ARID domain-containing polypeptides, such as the ones described herein, and the polypeptide fragment of GenBank accession number AAD41239 (PID g5257005).

5 By "E2F" or "E2F protein" is meant a protein having E2F transcriptional activity. Preferred E2F proteins include E2F-1, E2F-2, E2F-3, E2F-4, and E2F-5.

By "RB family member" is meant the tumor suppressors pRb, p130, and p107.

10 By an "HDAC" is meant any histone deacetylase protein. Preferred HDACs are HDAC-1, HDAC-2, and HDAC-3.

By a "substantially purified" protein or antibody is meant a protein or antibody that has been separated from components that naturally accompany it. Typically, the protein is substantially purified when it is at least 60%, by  
15 weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, a polypeptide of the invention. A substantially purified protein of the invention may be obtained, for example, by extraction from a natural source (for  
20 example, a virus), by expression of a recombinant nucleic acid encoding such a protein, or by chemically synthesizing the protein. Purity can be measured by any appropriate method, for example, column chromatography, polyacrylamide gel electrophoresis, or by HPLC analysis.

A protein is substantially free of naturally associated components when  
25 it is separated from those contaminants which accompany it in its natural state. Thus, a protein which is chemically synthesized or produced in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components. Accordingly,

-6-

substantially purified polypeptides include those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

By "compound" or "candidate compound" is meant a chemical, be it naturally-occurring or artificially-derived, that is assayed for its ability to modulate an alteration in reporter gene activity or protein levels, by employing one of the assay methods described herein. Test compounds may include, for example, peptides, polypeptides, synthesized organic molecules, naturally occurring organic molecules, nucleic acid molecules, and components thereof.

By "assaying" is meant analyzing the effect of a treatment, be it chemical or physical, administered to whole animals or cells derived therefrom. The material being analyzed may be an animal, a cell, a lysate or extract derived from a cell, or a molecule derived from a cell. The analysis may be, for example, for the purpose of detecting altered gene expression, altered RNA stability, altered protein stability, altered protein levels, or altered protein biological activity. The means for analyzing may include, for example, antibody labeling, immunoprecipitation, phosphorylation assays, and methods known to those skilled in the art for detecting nucleic acids and polypeptides.

By "modulating" is meant changing, either by decrease or increase, biological activity.

By "a decrease" is meant a lowering in the level of biological activity, as measured by a lowering of (i) RBP1 protein, as measured by ELISA, (ii) RBP1 transcriptional repression of E2F-dependent transcription, as measured, for example, by reporter gene assay (*lacZ*/ $\beta$ -galactosidase, green fluorescent protein, luciferase, etc.), (iii) RBP1 mRNA levels, as measured by PCR relative to an internal control, for example, a "housekeeping" gene product such as  $\beta$ -actin or glyceraldehyde 3-phosphate dehydrogenase (GAPDH), (iv) the amount of HDACs (or any other protein that is part of a protein complex that contains an HDAC) that interact with RBP1, (v) the



-7-

amount of E2F protein that interacts with RBP1, and (vi) RBP1-mediated cell cycle arrest. In all cases, the lowering is preferably by 20%, more preferably by 40%, and even more preferably by 70%.

By an "increase" is meant a rise in the level of biological activity, as measured by an increase of: (i) RBP1 protein, as measured by ELISA, (ii) RBP1 transcriptional repression of E2F-dependent transcription, as measured, for example, by reporter gene assay (*lacZ*/ $\beta$ -galactosidase, green fluorescent protein, luciferase, etc.), (iii) RBP1 mRNA levels, as measured by PCR relative to an internal control, for example, a "housekeeping" gene product such as  $\beta$ -actin or glyceraldehyde 3-phosphate dehydrogenase (GAPDH), (iv) the amount of HDACs (or any other protein that is part of a protein complex that contains an HDAC) that interact with RBP1, (v) the amount of E2F protein that interacts with RBP1, and (vi) RBP1-mediated cell cycle arrest. Preferably, the increase is by 5% or more, more preferably by 15% or more, even more preferably by 2-fold, and most preferably by at least 3-fold.

By "promoter" is meant a minimal sequence sufficient to direct transcription. Also included in the invention are those regulatory elements that are sufficient to render promoter-dependent gene expression controllable for cell type-specific, tissue-specific, temporal-specific, or inducible by external signals or agents; such elements may be located in the 5' or 3' or intron sequence regions of the native gene.

By "operably linked" is meant that a gene and a promoter are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the promoter.

By "pharmaceutically acceptable" is meant that a carrier is substantially non-toxic to the treated mammal and retains the therapeutic properties of the compound with which it is administered. One exemplary pharmaceutically acceptable carrier is physiological saline solution. Other physiologically

-8-

acceptable carriers and their formulations are known to one skilled in the art and described, for example, in Remington: The Science and Practice of Pharmacy, (19th ed.) ed. A.R. Gennaro, 1995, Mack Publishing Company, Easton, PA.

5 By a "matched" sample is meant one derived from the same or similar cells or tissue, from an organism of the same species, to which it is being compared.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof.

10 Description of the Drawings

Fig. 1 is a photograph of a autoradiograph showing analysis of E2F complexes by EMSA. Components of the E2F complexes were identified by antibody "supershift" experiments as described herein. Lane 1: asynchronously growing cells; lanes 2 to 4: extracts from asynchronously growing cells incubated in the presence anti-pRB serum (lane 2), anti-p107 serum SD9 (lane 3), or anti-p130 serum C-20 (lane 4); lane 5: serum-starved cells; lanes 6 to 8: extracts from serum-starved cells incubated in the presence of anti-pRB (lane 6), anti-p107 (lane 7) or anti-p130 (lane 8) sera; lanes 9 to 12: asynchronously growing cells (lane 9), serum-starved cells (lane 10), or extracts from serum-starved cells incubated in the presence of anti-p130 antibody (lane 11) or anti-RBP1 serum (lane 12); lanes 9A to 12A: the same samples as in lanes 9 to 12 analyzed using a polyacrylamide:bis-acrylamide ratio of 19:1. The positions of p130-E2F, and pRB-containing E2F complexes (RB) and those of "supershifted" complexes (\*p130-E2F and \*RB) are as indicated.

25 Figs. 2A and 2B are schematic illustrations showing the effect of RBP1 on E2F-dependent transcription. Human C33A cells were transiently transfected with E2F1-Luc(Wt) (Fig. 2A), which contains firefly luciferase

-9-

linked to an E2F-1 specific promoter, or E2F1-Luc(mE2F) (Fig. 2B), which contained mutated E2F recognition sites. They were also co-transfected with increasing amount (from 1  $\mu$ g - 4  $\mu$ g) of plasmids pCMVRBP1-HA (human RBP126), pCMVRBP1 $\Delta$ LCE-HA (a deletion mutant lacking RBP1 residues 5 921-1056 including the Leu-X-Cys-X-Glu binding site), and, in some cases, plasmids expressing human pRB, p130, or the product of the 12S Ad5 E1A mRNA subcloned into pcDNA3 (Invitrogen, Carlsbad, CA) and expressed under the cytomegalovirus (CMV) promoter. Cells were harvested at 40 hours post-transfection and luciferase activity was measured as previously described 10 (Weintraub et al., Nature 375:812-815, 1995). Results are presented as a percentage of that obtained with reporter plasmid alone. Fig. 2C is a schematic illustration showing the effects of RBP1(Wt) and RBP1( $\Delta$ LCE) on pRB and p130 repressed E2F1(Wt) promoter. Cells were transfected as in Fig. 2A. Additional samples were assayed by co-transfecting plasmids expressing pRB 15 or p130 with increasing amount of plasmids pCMVRBP1-HA expressing RBP1(Wt) or pCMVRBP1 $\Delta$ LCE-HA expressing RBP1( $\Delta$ LCE) mutant. Measured luciferase activities are presented as a percentage of that obtained with reporter plasmid alone and are plotted against amount of pCMVRBP1-HA or pCMVRBP1( $\Delta$ LCE) plasmids transfected.

20 Fig. 3A is a schematic illustration showing inhibition of colony formation by RBP1. CV-1 cells were transfected with plasmids expressing neo and either wild-type RBP1 or a truncated form of RBP1 that lacks the LXCXE-RB binding motif. Cells were grown under drug selection for two weeks. Colonies were then counted and the calculated averages were compared 25 with the number of colonies obtained with cells expressing neo alone (pcDNA3). Fig. 3B is a series of photographs showing morphology of colonies resulting from the experiment described in Fig. 3A.

Fig. 4A is a schematic illustration showing repression of Gal4-dependent

-10-

transcription by RBP1. Fig. 4B is a schematic illustration showing RBP1 deletion mutants and a summary of CAT activity as a percent of control.

Figs. 5A and 5B are a series of photographs showing interactions of pRB and p107 with histone deacetylases. The locations of hypophosphorylated species of pRB and p107 are as indicated.

Figs. 6A - 6D show *in vivo* interaction of the small 'pocket' of pRB with histone deacetylases. cDNAs encoding Gal4DBD fused with the 'pocket' of pRB and those encoding Flag-tagged HDAC1, HDAC2, or HDAC3 were overexpressed in H1299 (Fig. 6A). Anti-Gal4DBD antibody was used to immunoprecipitate Gal4-tagged overexpressed proteins and co-immunoprecipitation of HDACs was then detected by western blotting. WC - whole cell extracts; IP - anti-Gal4 immunoprecipitated proteins. Binding studies similar to those described in Fig. 6A were also performed in 293T cells (Fig. 6B) and 293 cells (Fig. 6C). A modified binding experiment similar to that described above was performed in H1299 cells by incubating cell extracts with increasing amounts (0 to 20  $\mu$ g) of E1A protein (first exon containing the LXCXE motif) fused to GST (GST-E1A) (Fig. 6D). The positions of migration of Flag-tagged HDAC1 (open arrows), HDAC2 (closed arrow) and HDAC3 (arrowhead) are as indicated.

Figs. 7A - 7E are a series of photographs showing *in vivo* interaction of RBP1 with histone deacetylases. Binding studies similar to those described in Fig. 6 were performed in H1299 cells (Fig. 7A) or 293T cells (Fig. 7B) using Gal4-RBP1 instead of Gal4-pRB. Binding studies were also performed in 293T cells transfected with the RBP1-HA construct and Gal4-HDAC1, Gal4-HDAC2, and Gal4-HDAC3 constructs (Fig. 7C). Coimmunoprecipitation studies similar were done in both H1299 and 293T cells. In addition to the antisera described, antibody against pRB, p130, p107, and the HA-epitope were also used (Fig. 7D). A binding study was also done using the Gal4-RBP1-

-11-

$\Delta$ LCE and RBP1- $\Delta$ LCE-HA mutants, each of which lacks the LXCXE 'pocket'-binding motif (Fig. 7E). The positions of migration of Flag-tagged HDAC1 (open arrows), HDAC2 (closed arrow) and HDAC3 (arrowhead) are as indicated.

5 Figs. 8A - 8D are a series of schematic illustrations showing mapping of transcriptional repression domains in RBP1 and the effect of trichostatin A (TSA) on RBP1 repression activity. CAT assays were performed in CHO cells using G5TKCAT as reporter. A summary of repression results obtained in experiments described in Fig. 8A are shown at the left of Fig. 8B. The effect of  
10 TSA on repression is shown in Fig. 8C. Repression assays were carried out as in Fig. 8A, except that some cells were incubated with 330 nM TSA for 24 hours prior to harvesting and the G5MLPCAT reporter was assayed instead of the G5TKCAT reporter. Fig. 8D shows a similar experiment to that described in Fig. 8C, except performed using a reporter construct consisting of the E2F-1  
15 promoter linked to a cDNA encoding luciferase and either RBP1, RBP1- $\Delta$ LCE, the 'pocket' of pRB, or an inactive pRB 'pocket'- mRB(C706F) mutant.

Fig. 9 is a schematic illustration showing mapping of specific binding of HDAC3 to RBP1. Binding studies were performed in 293T cells using either Gal4-R1, Gal4-R2(1314-C)/(1263-C), Gal4 alone, Gal4-RBP1(wt),  
20 Gal4-ARID, Gal4- $\Delta$ R1, or Gal4- $\Delta$ 93C( $\Delta$ -R2).

#### Detailed Description of the Invention

We have discovered that RBP1, a known pRB pocket-binding protein, possesses transcriptional repression activity and associates with p130-E2F and pRB-E2F complexes specifically during growth arrest. Overexpression of  
25 RBP1 both inhibited E2F-dependent gene expression and suppressed cell growth. Thus, we believe that repression of E2F-dependent transcription by RBP1 via RB family members plays a central role in inducing growth arrest.

-12-

We have also discovered that RBP1 interacts with histone deacetylases (HDACs). Unlike pRB-histone deacetylase interactions, however, 'pocket'-binding E1A protein and large T antigen do not affect interactions between RBP1 and any of the histone deacetylases. It was previously thought that HDAC1 or HDAC2 interacted directly with the RB pocket via degenerate IXCXE motifs, but this model does not account for interactions of HDAC3, which lacks such sequence. Our data indicated that binding of HDAC3 to RB family members also requires the pocket, suggesting that such interactions may be indirect and rely on a pocket-binding protein, such as RBP1. Our results show RBP1's functional importance in the RB family-mediated repression of E2F-dependent transcription, by linking histone deacetylases to the pockets of RB family proteins.

We have also identified two repressor domains in RBP1. One of these domains is HDAC-dependent, while the other is HDAC-independent.

Thus, we have found for the first time that RBP1 is a key member of the RB-E2F complex, serving in part as a link between RB and HDACs, but also serving a second, HDAC-independent, function. This discovery allows for the development of drugs that modulate or mimic RBP1 biological activity or that modulate its expression. These drugs have use in the treatment of disorders characterized by abnormal proliferation or cell death.

#### Production of RBP1 polypeptide

For protein expression, eukaryotic and prokaryotic expression systems may be generated in which *RBP1* gene sequences are introduced into a plasmid or other vector, which is then used to transform living cells. Constructs in which the RBP1 cDNAs containing the entire open reading frames inserted in the correct orientation into an expression plasmid may be used for protein expression. Alternatively, portions of the *RBP1* gene sequences, including

-13-

wild-type or mutant *RBP1* sequences, may be inserted. Prokaryotic and eukaryotic expression systems allow various important functional domains of the RBP1 proteins to be recovered as fusion proteins and then used for binding, structural and functional studies and also for the generation of appropriate antibodies. Since RBP1 protein expression may induce cell cycle arrest in some cell types, it may be desirable to express the protein under the control of an inducible promoter.

Typical expression vectors contain promoters that direct the synthesis of large amounts of mRNA corresponding to the inserted *RBP1* nucleic acid in the plasmid bearing cells. They may also include eukaryotic or prokaryotic origin of replication sequences allowing for their autonomous replication within the host organism, sequences that encode genetic traits that allow vector-containing cells to be selected for in the presence of otherwise toxic drugs, and sequences that increase the efficiency with which the synthesized mRNA is translated. Stable long-term vectors may be maintained as freely replicating entities by using regulatory elements of, for example, viruses (e.g., the OriP sequences from the Epstein Barr Virus genome). Cell lines may also be produced which have integrated the vector into the genomic DNA, and in this manner the gene product is produced on a continuous basis.

Expression of foreign sequences in bacteria, such as *Escherichia coli*, requires the insertion of the *RBP1* nucleic acid sequence into a bacterial expression vector. This plasmid vector contains several elements required for the propagation of the plasmid in bacteria, and expression of inserted DNA of the plasmid by the plasmid-carrying bacteria. Propagation of only plasmid-bearing bacteria is achieved by introducing in the plasmid selectable marker-encoding sequences that allow plasmid-bearing bacteria to grow in the presence of otherwise toxic drugs. The plasmid also bears a transcriptional promoter capable of producing large amounts of mRNA from the cloned gene. Such



-14-

promoters may or may not be inducible promoters, which initiate transcription upon induction. The plasmid also preferably contains a polylinker to simplify insertion of the gene in the correct orientation within the vector. In a simple *E. coli* expression vector utilizing the lac promoter, the expression vector plasmid contains a fragment of the *E. coli* chromosome containing the lac promoter and the neighboring *lacZ* gene. In the presence of the lactose analog IPTG, RNA polymerase normally transcribes the *lacZ* gene, producing *lacZ* mRNA, which is translated into the encoded protein,  $\beta$ -galactosidase. The *lacZ* gene can be cut out of the expression vector with restriction endonucleases and replaced by a *RBP1* gene sequence, or fragment, fusion, or mutant thereof. When this resulting plasmid is transfected into *E. coli*, addition of IPTG and subsequent transcription from the lac promoter produces *RBP1* mRNA, which is translated into a RBP1 polypeptide.

Once the appropriate expression vectors containing a *RBP1* gene, or fragment, fusion, or mutant thereof, are constructed, they are introduced into an appropriate host cell by transformation techniques including, for example, calcium phosphate transfection, DEAE-dextran transfection, electroporation, microinjection, protoplast fusion, and liposome-mediated transfection, as described herein. The host cells that are transfected with the vectors of this invention may include (but are not limited to) *E. coli*, *Pseudomonas*, *Bacillus subtilis*, or other bacilli, other bacteria, yeast, fungi, insect (using, for example, baculoviral vectors for expression), mouse or other animal or human tissue cells. Mammalian cells can also be used to express the RBP1 protein using a vaccinia virus expression system described in Ausubel et al., 1997, Current Protocols in Molecular Biology, Wiley Interscience, New York.

*In vitro* expression of RBP1 proteins, fusions, polypeptide fragments, or mutants encoded by cloned DNA is also possible using the T7 late-promoter expression system. This system depends on the regulated expression of T7



-15-

RNA polymerase which is an enzyme encoded in the DNA of bacteriophage T7. The T7 RNA polymerase transcribes DNA beginning within a specific 23-bp promoter sequence called the T7 late promoter. Copies of the T7 late promoter are located at several sites on the T7 genome, but none is present in *E. coli* chromosomal DNA. As a result, in T7 infected cells, T7 RNA polymerase catalyzes transcription of viral genes but not of *E. coli* genes. In this expression system recombinant *E. coli* cells are first engineered to carry the gene encoding T7 RNA polymerase next to the lac promoter. In the presence of IPTG, these cells transcribe the T7 polymerase gene at a high rate and synthesize abundant amounts of T7 RNA polymerase. These cells are then transformed with plasmid vectors that carry a copy of the T7 late promoter protein. When IPTG is added to the culture medium containing these transformed *E. coli* cells, large amounts of T7 RNA polymerase are produced. The polymerase then binds to the T7 late promoter on the plasmid expression vectors, catalyzing transcription of the inserted cDNA at a high rate. Since each *E. coli* cell contains many copies of the expression vector, large amounts of mRNA corresponding to the cloned cDNA can be produced in this system and the resulting protein can be radioactively labeled. Plasmid vectors containing late promoters and the corresponding RNA polymerases from related bacteriophages such as T3, T5, and SP6 may also be used for *in vitro* production of proteins from cloned DNA. *E. coli* can also be used for expression using an M13 phage such as mGPI-2. Furthermore, vectors that contain phage lambda regulatory sequences, or vectors that direct the expression of fusion proteins, for example, a maltose-binding protein fusion protein or a glutathione-S-transferase fusion protein, also may be used for expression in *E. coli*.

Eukaryotic expression systems permit appropriate post-translational modifications to expressed proteins. Transient transfection of a eukaryotic expression plasmid allows the transient production of a RBP1 polypeptide by a

-16-

transfected host cell. RBP1 proteins may also be produced by a stably-transfected mammalian cell line. A number of vectors suitable for stable transfection of mammalian cells are available to the public (e.g., see Pouwels et al., Cloning Vectors: A Laboratory Manual, 1985, Supp. 1987 and VectorDB at  
5 URL address <http://vectordb.atcg.com/>), as are methods for constructing such cell lines (see e.g., Ausubel et al., *supra*). In one example, cDNA encoding a RBP1 protein, fusion, mutant, or polypeptide fragment is cloned into an expression vector that includes the dihydrofolate reductase (DHFR) gene. Integration of the plasmid and, therefore, integration of the RBP1-encoding  
10 gene into the host cell chromosome is selected for by inclusion of 0.01-300  $\mu$ M methotrexate in the cell culture medium (as described, Ausubel et al., *supra*). This dominant selection can be accomplished in most cell types. Recombinant protein expression can be increased by DHFR-mediated amplification of the transfected gene. Methods for selecting cell lines bearing gene amplifications  
15 are described in Ausubel et al., *supra*. These methods generally involve extended culture in medium containing gradually increasing levels of methotrexate. The most commonly used DHFR-containing expression vectors are pCVSEII-DHFR and pAdD26SV(A) (described in Ausubel et al., *supra*). The host cells described above or, preferably, a DHFR-deficient CHO cell line  
20 (e.g., CHO DHFR<sup>-</sup> cells, ATCC Accession No. CRL 9096) are among those most preferred for DHFR selection of a stably-transfected cell line or DHFR-mediated gene amplification.

Eukaryotic cell expression of RBP1 proteins allows the production of large amounts of normal and mutant proteins for isolation and purification, and  
25 the use of cells expressing RBP1 proteins as a functional assay system for antibodies generated against the protein. Eukaryotic cells expressing RBP1 proteins may also be used to test the effectiveness of pharmacological agents on RBP1 biological activity, or as means by which to study RBP1 proteins as

-17-

components of a transcriptional regulatory complex. Expression of RBP1 proteins, fusions, mutants, and polypeptide fragments in eukaryotic cells also enables the study of the function of the normal complete protein, specific portions of the protein, or of naturally occurring polymorphisms and artificially produced mutated proteins. The *RBP1* DNA sequences can be altered using procedures known in the art, such as restriction endonuclease digestion, DNA polymerase fill-in, exonuclease deletion, terminal deoxynucleotide transferase extension, ligation of synthetic or cloned DNA sequences and site-directed sequence alteration using specific oligonucleotides together with PCR.

Another preferred eukaryotic expression system is the baculovirus system using, for example, the vector pBacPAK9, which is available from Clontech (Palo Alto, CA). If desired, this system may be used in conjunction with other protein expression techniques, for example, the myc tag approach described by Evan et al. (*Mol. Cell Biol.* 5:3610-3616, 1985).

Once the recombinant protein is expressed, it can be isolated from the expressing cells by cell lysis followed by protein purification techniques, such as affinity chromatography. In this example, an anti-RBP1 antibody can be attached to a column and used to isolate the recombinant RBP1 proteins. Lysis and fractionation of RBP1 protein-harboring cells prior to affinity chromatography may be performed by standard methods (see, e.g., Ausubel et al., *supra*). Once isolated, the recombinant protein can, if desired, be purified further by e.g., by high performance liquid chromatography (HPLC; e.g., see Fisher, Laboratory Techniques In Biochemistry And Molecular Biology, Work and Burdon, Eds., Elsevier, 1980).

Polypeptides of the invention, particularly short RBP1 fragments and longer fragments of the N-terminus and C-terminus of the RBP1 protein, can also be produced by chemical synthesis (e.g., by the methods described in Solid Phase Peptide Synthesis, 2nd ed., 1984, The Pierce Chemical Co., Rockford,

-18-

IL). These general techniques of polypeptide expression and purification can also be used to produce and isolate useful RBP1 polypeptide fragments or analogs, as described herein.

Those skilled in the art of molecular biology will understand that a wide  
5 variety of expression systems may be used to produce the recombinant RBP1  
proteins. The precise host cell used is not critical to the invention. The RBP1  
proteins may be produced in a prokaryotic host (e.g., *E. coli*) or in a eukaryotic  
host (e.g., *S. cerevisiae*, insect cells such as Sf9 cells, or mammalian cells such  
as COS-1, NIH 3T3, or HeLa cells). These cells are commercially available  
10 from, for example, the American Type Culture Collection, Rockville, MD (see  
also F. Ausubel et al., *supra*). The method of transformation and the choice of  
expression vehicle (e.g., expression vector) will depend on the host system  
selected. Transformation and transfection methods are described, e.g., in  
Ausubel et al., *supra*, and expression vehicles may be chosen from those  
15 provided, e.g., in Pouwels et al., *supra*.

#### Testing for the presence of RBP1 biological activity

Identification of both biologically active and mutant forms of RBP1  
allows the study of RBP1 biological activity in regulation of cell proliferation.  
For example, administration of an RBP1 protein, or a polypeptide fragment  
20 thereof, can be used to inhibit E2F-regulated gene expression, as measured by  
cell-based and cell-free assays known in the art and described herein. An  
inhibitory amount of an RBP1 reagent (e.g., a compound that reduces the  
biological function of RBP1, and thus may promote cell proliferation, such as  
an RBP1 neutralizing antibody or antisense *RBP1* nucleic acid, an *RBP1*  
25 nucleic acid encoding a dominant-negative form of the RBP1 protein, or a  
compound which decreases RBP1 gene expression) may be similarly assessed.  
Such assays may be carried out in a cell which either expresses endogenous

-19-

RBP1 or a cell to which is introduced a heterologous amount of an RBP1 polypeptide or in a cell-free assay. Preferably, the cell is capable of undergoing RBP1-mediated cell cycle arrest. RBP1 biological activity or inhibition thereof may be assessed in these RBP1 expressing cells, whereby such RBP1-inducing  
5 or -inhibiting activity is evaluated based upon the level of RBP1 biological activity.

One preferred RBP1 biological activity is binding of RBP1 to an HDAC. It is understood that the binding may not be through direct contact between RBP1 and the HDAC, but may require additional proteins. Binding of RBP1 to  
10 an HDAC, or any other protein that is part of a protein complex that contains an HDAC (e.g., SAP30, SinA, or SinB) can be measured using any of a number of standard methods known to those in the art, such as the methods described herein.

Another preferred RBP1 biological activity is binding of RBP1 to an  
15 E2F protein. As described for the RBP1-HDAC interaction, the binding of RBP1 to an E2F protein may require additional proteins.

A related RBP1 biological activity is repression of E2F transcriptional activity by RBP1. It is likely that the repression domains of RBP1 are sites in which proteins bind to RBP1 and repress E2F transcriptional activity. E2F  
20 transcriptional activity can be measured, for example, using the methods described herein.

In another embodiment, RBP1 biological activity is slowing of the cell cycle or cell cycle arrest. There are numerous cell cycle assays known in the art, including those described herein; any of these may be used in measuring  
25 RBP1-mediated cell cycle regulation.

#### Identification of molecules that modulate RBP1 biological activity

Methods of observing changes in RBP1 biological activity, such as the

-20-

ones described above, are exploited in high-throughput assays for the purpose of identifying compounds that modulate mutant or wild-type RBP1 transcriptional activity. Compounds that mime RBP1 activity also may be identified by such assays. Furthermore, compounds that modulate transcription of the *RBP1* gene itself may be identified; in some cases, it may be desirable to increase or decrease RBP1 protein levels by such an approach. Such identified compounds may have utility as therapeutic agents in the treatment of cell proliferation disease.

In general, novel drugs that modulate or mime RBP1 biological activity or expression are identified from large libraries of both natural product or synthetic (or semi-synthetic) extracts or chemical libraries according to methods known in the art. Those skilled in the field of drug discovery and development will understand that the precise source of test extracts or compounds is not critical to the screening procedure(s) of the invention. Accordingly, virtually any number of chemical extracts or compounds can be screened using the exemplary methods described herein. Examples of such extracts or compounds include, but are not limited to, plant-, fungal-, prokaryotic- or animal-based extracts, fermentation broths, and synthetic compounds, as well as modification of existing compounds. Numerous methods are also available for generating random or directed synthesis (e.g., semi-synthesis or total synthesis) of any number of chemical compounds, including, but not limited to, saccharide-, lipid-, peptide-, and nucleic acid-based compounds. Synthetic compound libraries are commercially available from Brandon Associates (Merrimack, NH) and Aldrich Chemical (Milwaukee, WI). Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant, and animal extracts are commercially available from a number of sources, including Biotics (Sussex, UK), Xenova (Slough, UK), Harbor Branch Oceanographics Institute (Ft. Pierce, FL), and PharmaMar, U.S.A. (Cambridge,

-21-

MA). In addition, natural and synthetically produced libraries are produced, if desired, according to methods known in the art, e.g., by standard extraction and fractionation methods. Furthermore, if desired, any library or compound is readily modified using standard chemical, physical, or biochemical methods.

5 In addition, those skilled in the art of drug discovery and development readily understand that methods for dereplication (e.g., taxonomic dereplication, biological dereplication, and chemical dereplication, or any combination thereof) or the elimination of replicates or repeats of materials already known for their therapeutic activities for cell proliferation diseases  
10 should be employed whenever possible.

When a crude extract is found to modulate RBP1 biological activity, further fractionation of the positive lead extract is necessary to isolate chemical constituents responsible for the observed effect. Thus, the goal of the extraction, fractionation, and purification process is the careful characterization  
15 and identification of a chemical entity within the crude extract that modulates or mimics RBP1 biological activity or expression. The same assays described herein for the detection of activities in mixtures of compounds can be used to purify the active component and to test derivatives thereof. Methods of fractionation and purification of such heterogenous extracts are known in the  
20 art. If desired, compounds shown to be useful agents for treatment are chemically modified according to methods known in the art. Compounds identified as being of therapeutic value may be subsequently analyzed using a standard animal or culture model for a cell proliferative disease known in the art.

25 The effect of candidate compounds on RBP1-mediated regulation of cell growth or survival may be measured at the level of translation by using the general approach described above with standard protein detection techniques, such as Western blotting or immunoprecipitation with an RBP1-specific



-22-

antibody (for example, the RBP1-specific antibody described herein).

Compounds that modulate the level of RBP1 may be purified, or substantially purified, or may be one component of a mixture of compounds such as an extract or supernatant obtained from cells (Ausubel et al., *supra*). In  
5 an assay of a mixture of compounds, RBP1 expression is measured in cells administered progressively smaller subsets of the compound pool (e.g., produced by standard purification techniques such as HPLC or FPLC) until a single compound or minimal number of effective compounds is demonstrated to modulate RBP1 expression.

10 Compounds may also be screened for their ability to modulate RBP1 biological activity. In this approach, the degree of RBP1 biological activity in the presence of a candidate compound is compared to the degree of biological activity in its absence, under equivalent conditions. Again, the screen may begin with a pool of candidate compounds, from which one or more useful  
15 modulator compounds are isolated in a step-wise fashion. Biological activity may be measured by any standard assay, for example, those described herein.

Another method for detecting compounds that modulate the activity of RBP1 is to screen for compounds that bind to RBP1. These compounds may be detected by adapting interaction trap expression systems known in the art.  
20 These systems detect protein interactions using a transcriptional activation assay and are generally described by Gyuris et al. (Cell 75:791-803, 1993) and Field et al., (Nature 340:245-246, 1989), and are commercially available from Clontech (Palo Alto, CA). Alternatively, compounds that bind to RBP1 can be isolated by column chromatography. In this assay, candidate compounds or  
25 extracts are passed through a column containing immobilized RBP1. After washing to remove nonspecific proteins, the compounds bound to RBP1 are eluted, collected, and identified. Using similar assays, compounds that interfere with RBP1 binding to pRB family members or HDACs can also be



-23-

identified.

Compounds or molecules that modulate RBP1 biological activity may include peptide and non-peptide molecules such as those present in cell extracts, mammalian serum, or growth medium in which mammalian cells have  
5 been cultured.

#### Therapeutic uses

A molecule that decreases RBP1 expression or biological activity is considered useful in the invention; such a molecule may be used, for example, as a therapeutic to increase cell proliferation in an arrested cell (e.g., a neuron).  
10 There is a tight association between cell proliferation and cell death. In some cases, a cell that receives a signal to proliferate will instead undergo apoptosis, presumably because the proliferation is inappropriate. Hence, depending on the context, a molecule that decreases RBP1 expression or biological activity may result in increased proliferation or, alternatively, cell death. Differentiating  
15 between these two outcomes in the appropriate context may readily be done using standard assays known to those skilled in the art.

A molecule that increases RBP1 expression or biological activity is particularly useful in the invention. Such a molecule may be used to decrease cellular proliferation. This would be advantageous in the treatment of  
20 neoplasms or other cell proliferative diseases.

To add an RBP1 polypeptide to cells in order to modulate cell proliferation or apoptosis, it is preferable to obtain pure RBP1 protein from cultured cell systems that can express the protein. Delivery of the protein to the affected tissue can then be accomplished using appropriate packaging or  
25 administering systems. Alternatively, small molecule analogs may be used and administered to act as RBP1 agonists or antagonists and in this manner produce a desired physiological effect. Methods for finding such molecules are

-24-

provided herein.

Gene therapy is another potential therapeutic approach in which normal copies of the RBP1 gene or nucleic acid encoding RBP1 sense or antisense RNA is introduced into cells. The gene must be delivered to those cells in a form in which it can be taken up and encode for sufficient protein to provide effective function.

Transducing retroviral, adenoviral, and human immunodeficiency viral (HIV) vectors can be used for somatic cell gene therapy especially because of their high efficiency of infection and stable integration and expression (see, for example, Cayouette and Gravel, Hum. Gene Ther., 8:423-430, 1997; Kido et al. Curr. Eye Res., 15:833-844, 1996; Bloomer et al., J. Virol., 71:6641-6649, 1997; Naldini et al., Science 272:263-267, 1996; Miyoshi et al., Proc. Natl. Acad. Sci. USA, 94:10319-10323, 1997). For example, the full length RBP1 nucleic acid, or portions thereof, can be cloned into a retroviral vector and driven from its endogenous promoter or from the retroviral long terminal repeat or from a promoter specific for the target cell type of interest (such as neurons). Other viral vectors which can be used include adenovirus, adeno-associated virus, vaccinia virus, bovine papilloma virus, or a herpes virus, such as Epstein-Barr Virus.

Gene transfer could also be achieved using non-viral means requiring infection *in vitro*. This would include calcium phosphate, DEAE dextran, electroporation, and protoplast fusion. Liposomes may also be potentially beneficial for delivery of DNA into a cell. Although these methods are available, many of these are of lower efficiency.

Transplantation of normal genes into the affected cells of a patient can also be useful therapy. In this procedure, a normal RBP1 gene is transferred into a cultivatable cell type, either exogenously or endogenously to the patient. These cells are then injected into the targeted tissue(s).

-25-

In the constructs described, RBP1 cDNA expression can be directed from any suitable promoter (e.g., the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element. For example, if desired, enhancers  
5 known to preferentially direct gene expression in tumor cells may be used to direct RBP1 expression. The enhancers used could include, without limitation, those that are characterized as tissue- or cell-specific in their expression. Alternatively, if a RBP1 genomic clone is used as a therapeutic construct (for example, following isolation by hybridization with the RBP1 cDNA described  
10 above), regulation may be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

Antisense based strategies may be employed to explore RBP1 gene function and as a basis for therapeutic drug design. The principle is based on  
15 the hypothesis that sequence-specific suppression of gene expression can be achieved by intracellular hybridization between mRNA and a complementary antisense species. The formation of a hybrid RNA duplex may then interfere with the processing/transport/translation and/or stability of the target RBP1 mRNA. Antisense strategies may use a variety of approaches including the use  
20 of antisense oligonucleotides and injection of antisense RNA. Antisense effects can be induced by control (sense) sequences, however, the extent of phenotypic changes are highly variable. Phenotypic effects induced by antisense effects are based on changes in criteria such as protein levels, protein activity measurement, and target mRNA levels.

25 For example, RBP1 gene therapy may also be accomplished by direct administration of antisense *RBP1* mRNA to a cell in which proliferation is desired. The antisense *RBP1* mRNA may be produced and isolated by any standard technique, but is most readily produced by *in vitro* transcription using

-26-

an antisense *RBP1* cDNA under the control of a high efficiency promoter (e.g., the T7 promoter). Administration of antisense *RBP1* mRNA to cells can be carried out by any of the methods for direct nucleic acid administration described above.

- 5           An alternative strategy for inhibiting RBP1 function using gene therapy involves intracellular expression of an anti-RBP1 antibody or a portion of an anti-RBP1 antibody. For example, a nucleic acid (or fragment thereof) encoding a monoclonal antibody that specifically binds to RBP1 and inhibits its biological activity may be placed under the transcriptional control of a cell  
10   type-specific gene regulatory sequence.

- Another therapeutic approach within the invention involves administration of a recombinant RBP1 polypeptide (e.g, the one described herein), either directly to the site of a potential or actual cell proliferation event (for example, by injection) or systemically (for example, by any conventional  
15   recombinant protein administration technique). The dosage of RBP1 depends on a number of factors, including the size and health of the individual patient, but, generally, between 0.1 mg and 100 mg inclusive are administered per day to an adult in any pharmaceutically acceptable formulation.

Administration of RBP1 Polypeptides, *RBP1* Genes, or Modulators of RBP1  
20   Synthesis or Function

- An RBP1 protein, gene, or modulator of RBP1 may be administered within a pharmaceutically-acceptable diluent, carrier, or excipient, in unit dosage form. Conventional pharmaceutical practice may be employed to provide suitable formulations or compositions to administer activating RBP1  
25   antibodies or compounds that activate or mimic RBP1 biological activity to patients suffering from a cell proliferation disease. Administration may begin before the patient is symptomatic. Methods well known in the art for making

-27-

- formulations are found, for example, in Remington: The Science and Practice of Pharmacy (*supra*). Formulations for parenteral administration may, for example, contain excipients, sterile water, or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes.
- 5 Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of the compounds. Other potentially useful parenteral delivery systems for RBP1 modulatory compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes.
- 10 Formulations for inhalation may contain excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel.

#### Gene Regulation

- 15 The transcriptional repressor domains described herein can be used to repress expression of any gene by targeting them to the regulatory region of that gene; the presence of the repressor domain will block transcriptional activation (Badiani et al., Genes Dev. 8:770-782, 1994; Conlon et al., Development 122:2427-2435, 1996). Targeting of the repressor domain can be
- 20 achieved by producing a fusion protein containing the repressor domain fused in frame to a DNA binding domain known to bind to the desired gene regulatory region. The fusion protein can be produced using standard methods and administered in a pharmaceutically-acceptable carrier to the cell or tissue of interest as described herein for the RBP1 polypeptides. Alternatively, the
- 25 nucleic acid sequence encoding the fusion protein can be used in gene therapy, as described above.

The following examples are to illustrate the invention. They are not

-28-

meant to limit the invention in any way.

Example 1: RBP1 is present in E2F-RB family complexes

Because of its slow migration in E2F-specific electrophoretic mobility shift assays (EMSA), we hypothesized that the p130-E2F complex appeared to  
5 contain one or more additional proteins and that such proteins interact with p130 via a site in the RB pocket similar to that utilized by the adenovirus E1A protein which binds via a conserved Leu-X-Cys-X-Glu motif found in several other viral and cellular RB-binding proteins.

To determine if RBP1 was present in E2F complexes, we harvested  
10 monkey CV-1 cells from either rapidly growing or serum-starved growth-arrested cultures, and subjected cell extracts to electrophoretic mobility shift assays (EMSA) using an E2F-specific radioactive oligonucleotide, as described previously (Corbeil et al., Oncogene 15:657-668, 1997). Fig. 1 shows that serum-starved (lanes 5 and 10) and asynchronously growing cells  
15 (lanes 1 and 9) contain a variety of E2F complexes; we observed the large p130-E2F complex only in the former. No complexes were detected using an unrelated or altered E2F oligonucleotide. The components of these E2F complexes were partially resolved through the addition of antibodies that recognize various RB family members. Addition of anti-pRB serum caused a  
20 supershift of at least two faster migrating pRB-E2F complexes (termed RB in the figure) to slower migrating species (\*RB) in both growing (lane 2) and serum-starved (lane 6) cells. We did not observe any effect on the migration of p130-E2F with this serum (lane 6). Addition of anti-p107 antibody SD9 had little effect on any of the complexes (lanes 3 and 7); addition of C-20 serum  
25 specific for the carboxyl terminus of p130, however, caused a supershift of p130-E2F in extracts from growth-arrested cells (lanes 8 and 11). Anti-p130 serum also made evident the existence of more minor, faster-migrating

-29-

p130-E2F complexes that were evident in lane 11, upon prolonged exposure of the gel. To determine if RBP1 is present in p130-E2F or any of the other E2F complexes, we conducted a similar experiment in asynchronously growing and serum-starved CV-1 cells using antiserum against human RBP1. Fig. 1 (lane 5 12) shows that both p130-E2F and the faster-migrating pRB-E2F complexes, identified in lane 2, had disappeared and that one or more new complexes (termed \*RB in the figure) were evident. In addition, analysis of these same samples under different conditions of PAGE indicated the presence of an additional, very slowly-migrating E2F complex (termed p130-E2F in Fig. 1) 10 that just barely entered the gel (lane 12A). Thus it appeared that p130-E2F and certain pRB-E2F complexes in growth arrested cells contain RBP1.

#### Example 2: RBP1 inhibits E2F-mediated transcription

To analyze the role of RBP1 in RB-E2F complexes, we analyzed rapidly growing C33A human cervical carcinoma cells, which are deficient both for 15 functional p53 and pRB, for their ability to regulate the E2F-1 promoter. We chose this promoter because it contains E2F recognition sites that are tightly regulated during growth arrest by E2F-RB family complexes, particularly p130/E2F (Hsiao et al., Genes Dev. 8:1526-1537, 1994; Johnson et al., Nature 365:349-352, 1993; Johnson et al., Genes Dev. 8, 1514-1525, 1994; Johnson, 20 Oncogene 11:1685-1692, 1995). We transfected cells with DNA from E2F1-Luc(Wt), a luciferase reporter construct driven by the E2F-1 promoter, together with pCMVRBP1-HA plasmid that encodes a haemagglutinin (HA) epitope-tagged RBP1, or pCMVRBP1 $\Delta$ LCE-HA (a mutant RBP1 lacking the Leu-X-Cys-X-Glu p130/pRB binding site). In some cases, we also assayed 25 DNA from plasmids encoding pRB, p130, or E1A-243R (the 243-residue product of human adenovirus type 5 12S E1A mRNA). As shown in Fig. 2A, increasing the amount of cotransfected RBP1(Wt) resulted in decreasing



-30-

amount of luciferase activity. Thus, expression is comparable to the transcriptional repression effect that p130 and pRB each exerted on this E2F1-Luc(wt) reporter. Overexpression of RBP1 caused about a 50% reduction in luciferase activity, whereas introduction of the  $\Delta$ LCE mutant had no inhibitory effect. Expression of E1A-243R increased luciferase activity by almost nineteen-fold, as was expected. To determine if RBP1-mediated transcriptional repression was specific to the E2F-regulated transcription on this promoter, we used an E2F1-luc(mE2F) reporter, which contains four point mutations within the E2F recognition sites (Johnson, Oncogene 11:1685-1692, 1995). This mutated sequence was unable to interact with E2F in EMSA (Johnson et al., Genes Dev. 8, 1514-1525, 1994). As shown in Fig. 2B, the relative luciferase activity of this reporter was about three- to four-fold higher than the E2F1-luc(wt) reporter. Overexpression of either RBP1(Wt) or RBP1( $\Delta$ LCE) had no effect on the mutated E2F1 promoter, suggesting the transcriptional repression by RBP1 is specific to the E2F element within the E2F1 promoter. Taken with the finding that RBP1 and pRB are together in a complex, these results suggest that RBP1 inhibits E2F-dependent transcription and that such inhibition requires binding to RB family members. We obtained similar results with other cell types, including CHO C33A cells, which lack functional pRB. Thus, endogenous p130 and/or p107 must be sufficient for this inhibition.

Overexpression of either p130 or pRB has no effect on the E2F1-luc(mE2F) promoter, whereas overexpression of E1A-243R results in about three-fold activation of this promoter (Fig. 2B). E1A has been shown to enhance Sp1 driven transcription (Weintraub and Dean, Nature 358:259-261, 1992). Additionally, both the wild-type and mutated E2F-1 promoters contain a series of Sp1 recognition sites. These data explain E1A-enhanced transcription of both promoters, as shown in Figs. 2A and 2B. The fact that the



-31-

E2F1-luc(Wt) promoter was activated by E1A stronger than the E2F1-luc(mE2F) promoter, suggests that the E2F element within the E2F1 promoter is also effected by E1A overexpression.

5 Example 3: RBP1 inhibition of E2F-mediated transcription is dependent on interaction with RB family members

We predicted that overexpression of RBP1, together with RB family members, should cause further inhibition to the E2F-1 promoter. This was observed, as shown in Fig. 2C. With increasing amount of RBP1(Wt), along with constant amount of cotransfected p130 or pRB, we detected further inhibition of luciferase activity. Interestingly, with increasing amount of RBP1( $\Delta$ LCE), along with constant amount of cotransfected p130 or pRB, we observed a gradual increase of luciferase activity. Our observation of de-repression by the RBP1( $\Delta$ LCE) mutant suggests one or more common factor(s), required for RBP1 transcriptional repression effect on the E2F1 promoter, interact with both endogenous RBP1 and transfected RBP1( $\Delta$ LCE). Overexpression of RBP1( $\Delta$ LCE) would sequester these essential factors from the E2F-1 promoter due to the inability of  $\Delta$ LCE mutant to interact with the RB family members. The inhibitory effect on E2F-1 promoter by RBP1 required the interaction with RB family members through the LXCXE motif.

20 Accordingly RBP1( $\Delta$ LCE) is likely to act as a dominant negative mutant, partially abolishing the transcriptional repression effect of exogenously transfected p130 or pRB on the E2F-1 promoter (Fig. 2C). The fact that RBP1( $\Delta$ LCE) could only partially abolish the transcriptional repression effect exerted by the cotransfected p130 or pRB, suggests that pRB or p130 repress

25 transcription by other mechanisms, possibly through interaction with other LXCXE-containing cellular proteins.

-32-

Example 4: Overexpression of RBP1 induces growth arrest

As the E2F-p107 complex containing RBP1 is only formed during growth arrest, it seemed possible that RBP1 might play a role in the control of cell proliferation by inhibiting E2F-dependent transcription, thus inducing, enhancing, or maintaining exit from the cell cycle. To determine if RBP1 suppressed cell growth, we transfected CV-1 cells with DNA from plasmids that provide neomycin resistance and express RBP1. Constructs included pRcCMVRBP1-HA (encoding an HA-tagged version of RBP1), pcDNA3Gal4-RBP1 (containing RBP1 fused to the Gal4 DNA-binding motif); and pcDNA3Gal4-RBP1 $\Delta$ LCE. Following transfection, we plated cells and allowed them to grow under G418 selection. Fig. 3A shows that no colonies were formed in the absence of vectors. With pcDNA3Gal4-RBP1 $\Delta$ LCE, a significant number of large colonies were produced with a morphology similar to those obtained with the control pcDNA3, lacking RBP1 coding sequences (Fig. 3B). With each of the plasmids expressing wild-type RBP1, we observed a significant reduction in colony formation. In addition, the colonies of cells transfected wild-type RBP1 were considerably smaller and composed of only small numbers of sparse cells (see Fig. 3B).

To examine the mechanism by which RBP1 inhibits E2F activity, we fused the GAL4 DNA binding sequence to RBP1. We then co-transfected CHO cells with DNA encoding GAL4-RBP1 (or control proteins) and G5TKCAT, a reporter construct that expresses CAT by GAL4-dependent transcription. Fig. 4A shows that GAL4-RBP1, expressed from either the CMV or SV40 promoter, repressed transcription by 80 to 90%. This expression was as great as that by the adenovirus E1B-55kDa product (Gal4AdE1B-55K), known to be a potent transcriptional repressor when linked to the Gal4 DNA binding domain. RBP1-HA that was not linked to Gal4 (pRcCMVRBP1-HA) had no effect on expression of CAT. Thus, RBP1

-33-

appears to function as a transcriptional repressor, but only when tethered to a promoter. GAL4-RBP1 $\Delta$ LCE also exhibited similar repression activity, indicating that repression by RBP1 is independent of RB family binding when fused to a heterologous DNA binding domain.

5 To identify the region of RBP1 required for repression, we created a series of mutants in the GAL4-RBP1 construct in which increasing regions from the carboxyl terminus of RBP1 were deleted (see Fig 4B). Fig. 4 shows that elimination of sequences encoding up to 805 residues from the carboxyl terminus resulted in little or no reduction of repression activity; removal of  
10 additional sequences, however, greatly reduced or eliminated repression. It is likely that residues 241 to 452 are responsible for RBP1 repression activity. This portion of RBP1 contains a sequence predicted to encode an  $\alpha$ -helical domain, preceded by a region, termed ARID, found in several other proteins, including the product of dead ringer (dri) from *Drosophila melanogaster*  
15 (Gregory et al., Mol. Cell. Biol. 16:792-799, 1996), SWI1/ADR6 from yeast (Coté et al., Science 265:53-60, 1994), B Cell Regulator of IgH Transcription from mouse (Herrscher et al., Genes Dev. 9:3067-3082, 1995), human Mrf1 and Mrf2A (Huang et al., Nucl. Acids Res. 24:1695-1701, 1996), and RBP2 (Defeo-Jones et al., Nature 353:251-254, 1991; Fattaey et al., Oncogene  
20 8:3149-3156, 1993; Kaelin et al., Cell 70: 351-364, 1992; Kim et al., Mol. Cell. Biol. 14: 7256-7264, 1994; Otterson et al., Oncogene 8:949-957, 1993). When we fused residues 241 to 452 of RBP1 to Gal4 (Gal4-RBP1-R1), the resulting fusion protein repressed transcription at a level comparable to wild type Gal4-RBP1 (Fig. 4A), suggesting that this region is responsible for the  
25 transcriptional repression activity.

Example 5: Endogenous interactions between RB family members and histone deacetylases

-34-

We conducted co-immunoprecipitation experiments to determine the ability of pRB to interact with various HDAC enzymes *in vivo*. Rabbit polyclonal antibodies that specifically recognize HDAC2, or HDAC1 and HDAC3 were used under low stringency conditions to immunoprecipitate endogenous HDAC species from extracts of human H1299 cells. In each case, we immunoprecipitated significant amounts of corresponding HDAC proteins. We determined the presence of pRB in these complexes by Western blot analysis using G3-245 monoclonal antibody that recognizes both hyperphosphorylated and hypophosphorylated forms of pRB. Fig. 5A shows that the hypophosphorylated form of pRB co-precipitated with HDAC1. Interestingly, we also detected significant amounts of hypophosphorylated pRB in association with HDAC2 and HDAC3. These interactions were specific as we did not detect pRB in immunoprecipitates prepared from the same cell extracts using antibody to CREB-binding protein (CBP), a histone acetyltransferase. These results indicate that the active form of pRB is able to recruit all three forms of HDACs.

HDAC1 also interacts with pRB-related family members p107 and p130. We immunoblotted immunoprecipitates from H1299 cells containing various HDACs with C-18 polyclonal antibody, which recognizes up to three different phosphorylated forms of p107 in various cell lines. Fig. 5B shows that p107, in particular its underphosphorylated forms, also co-precipitated with all endogenous HDAC species. We attempted a similar experiment with p130, but, as H1299 cells die upon serum starvation and p130 is expressed largely at growth arrest, we were unable to detect sufficient quantities. Significant amounts of p130 were present in asynchronous H1299 cells, but these p130 species are mostly hyperphosphorylated and did not appear to associate with HDACs at significant levels.

Adenovirus E1A protein and SV40 large T antigen associate with the

-35-

pRB pocket via LXCXE binding motifs and disrupt interactions between pRB and HDAC1. We therefore conducted a parallel series of binding studies in 293T cells, which express high levels of both Ad5 E1A proteins and SV40 large T antigen. Fig. 5A shows that no interactions were apparent between pRB and any of the HDAC enzymes, including HDAC3 in these cells. Fig. 5B shows that a similar effect was apparent with p107. Thus in both cases, binding of all three HDAC enzymes seemed to require the RB pocket region targeted by DNA tumor virus proteins.

Example 6: The small pocket of pRB interacts with different histone deacetylases

It has been proposed that HDAC1 utilizes a degenerate IXCXE motif to interact with the pocket (residues 379 to 792) of pRB (Ferreira et al., Proc. Natl. Acad. Sci. USA 95:10493-10498, 1998; Magnaghi-Jaulin et al., Nature 391:601-605, 1998). To determine directly if this region is involved in binding of all of the HDACs, we carried out studies using extracts from H1299 cells co-transfected with plasmid DNA expressing the small pocket of pRB as a fusion product with the DNA binding domain of Gal4 (Gal4DBD), termed Gal4-RB(pocket), as well as Flag-tagged versions of HDAC1, 2, or 3. Following immunoprecipitation using an antibody against Gal4DBD, we resolved precipitates by SDS-PAGE and, after transfer, analyzed the presence of HDAC1, 2, or 3 by Western blotting using anti-Flag antibody. Fig. 6A shows that all three HDACs, which were detected at similar levels in whole cell extracts, associated with the small pocket of pRB *in vivo*. Anti-Flag antibody recognized three Flag-HDAC1 species, but only the slowest migrating form was evident in Gal4-RB(pocket) precipitates, suggesting that the faster migrating species may be degradation products. We detected Flag-HDAC3 as two closely migrating species and both associated with the Gal4-RB(pocket).

-36-

These interactions were highly specific as a cDNA expressing Gal4DBD linked to VP16, a known transcriptional activator, did not associate with any of the histone deacetylases (Fig. 6A) even though Gal4-RB(pocket) and Gal4-VP16 were expressed at comparable levels. These results, as well as those described above with anti-CBP antibody, demonstrate clearly the specificity of interactions involving HDACs, and rule out the possibility that the interactions we observed were non-specific.

Fig. 6B shows that in a similar experiment in 293T cells, none of the HDAC proteins were present in immunoprecipitates containing Gal4-RB(pocket). Identical results were also obtained in 293 cells that express only Ad5 E1A proteins (Fig. 6C). Fig. 6D shows that addition prior to immunoprecipitation of increasing amounts of *in vitro* synthesized E1A protein fused to GST (GST-E1A) (Corbeil et al., J. Virol. 68:6697-6709, 1994) to extracts from H1299 cells expressing Gal4-pRB(pocket) and Flag-HDAC1 caused a decrease in binding of HDAC1 to the small pocket of pRB. We did not observe such an effect with GST/glutathione-Sepharose alone. Fig. 6D also demonstrates that pRB coimmunoprecipitated with E1A protein but not with GST. We obtained similar results using Flag-HDAC2 and Flag-HDAC3. Thus the small pocket of pRB is important for interactions not only with HDAC1, but also with HDAC2 and HDAC3. Disruption of such binding by E1A protein or T antigen also implied that the association of histone deacetylases may be mediated by LXCXE-like interactions. Although HDAC1 and HDAC2 contain LXCXE motifs, the mechanism of binding of HDAC3 was uncertain as HDAC3 lacks such sequences. Of further interest, no interaction was previously observed between the pocket of pRB and HDAC1 or HDAC2 in studies involving the yeast two-hybrid method (Brehm et al., Nature 391:597-601, 1998). As all three human histone deacetylases are highly homologous, it was possible that interactions of all three with the pocket may be indirect, and

-37-

involve an additional LXCXE-containing protein as a linker. One possible candidate is RBP1.

Example 7: RBP1 interacts with histone deacetylases

We tested the ability of RBP1 to associate with histone deacetylases.

- 5 Fig. 7A shows that, when expressed in H1299 cells, Gal4-RBP1 (but not Gal4-VP16) interacted with all three histone deacetylases. Again, only the slowest migrating Flag-HDAC1 species was evident and Flag-HDAC3 was present as a doublet in Gal4-RBP1 precipitates. Fig. 7B shows that in 293T cells expressing E1A products and SV40 large T antigen, both HDAC1 and
- 10 HDAC3 associated with Gal4RBP1, indicating that these interactions were unaffected by high levels of these viral pocket-binding proteins, and thus did not require the binding to RB. Although binding of HDAC2 to RBP1 was not evident in Fig. 7B, residual amounts of HDAC2 were detected after longer exposures of the blot. In addition, we observed a low level of binding of
- 15 HDAC2 to RBP1 in a parallel experiment, shown in Fig. 6C, using 293 cells. Flag-HDAC2 expression was consistently low in 293T or 293 cells and this effect probably resulted from the fact that Flag-HDAC2 was expressed under a hybrid SV40/HTLVII promoter, whereas Flag-HDAC1 and Flag-HDAC3 expression relied on the CMV promoter. Thus, we re-examined RBP1-HDAC
- 20 binding in an additional experiment in which all three HDACs were expressed along with HA-tagged RBP1 in 293T cells as Gal4DBD-HDAC fusion products under the SV40 early promoter. Fig. 7C shows that all HDACs were expressed at comparable levels as determined by immunoblotting of whole cell extracts with anti-Gal4DBD antibody. Similarly, all cells expressed
- 25 comparable amounts of RBP1, as detected using anti-HA antibody. Fig. 7C



-38-

with anti-HA antibody. We observed no such binding with Gal4 alone or with Gal4-VP16. Fig. 7D shows that interactions between RBP1 and endogenous HDAC enzymes could also be detected in H1299 cells. With immunoprecipitates prepared using the same polyclonal antibodies against individual HDACs employed in Figs. 5A and 5B, we found RBP1 in association with all three HDACs. RBP1 interactions with HDAC1 clearly occurred at much higher levels than with HDAC2 and HDAC3, implying that RBP1/HDAC1 complexes represent the predominant species in H1299 cells. This observation may suggest that RBP1 could be a component of the HDAC1 core complex. If this is the case, much of the endogenous RBP1/HDAC1 complex could be targeted to sites other than RB family members as the levels of RBP1 detected in association with these proteins were much lower than those observed with HDAC1. Fig. 7D also shows that RBP1 associates with all members of the RB family, and examination of pertinent lanes in Figs. 5A and 5B indicated that this association occurred preferentially with hypophosphorylated forms of pRB (Fig. 1A) and p107 (Fig. 1B). In addition, Fig. 7D shows that such RBP1-pRB/p107/p130 interactions were disrupted in 293T cells, as expected. Taken together, these results indicate that interactions between RBP1 and RB family members are truly pocket-dependent, as they were disrupted by T antigen or E1A pocket-binding proteins. To the contrary, interactions between RBP1 and HDACs were not sensitive to disruption by these viral pocket-binding proteins, suggesting that HDACs cannot be recruited to RBP1 via endogenous RB family members present in 293T cells. Instead, RBP1 could mediate the association between RB family members and all three HDAC enzymes.

It is unlikely that interactions of histone deacetylases with RBP1 occur indirectly through the recruitment of RB family members by RBP1. In 293T/293 cells, we failed to observe interactions either between the pocket of



-39-

pRB and these enzymes (Figs. 2B and 2C) or between RBP1 and RB family members (Figs. 1A, 1B, and 3D). In addition, we studied interactions between histone deacetylases and RBP1( $\Delta$ LCE). Fig. 7E shows that in 293T cells, both Gal4RBP1( $\Delta$ LCE) or RBP1( $\Delta$ LCE)-HA mutants interacted with Flag-tagged or Gal4DBD-tagged histone deacetylases. We obtained similar results in human H1299 cells. Thus, histone deacetylases appear to interact with RBP1 in a region apart from the pocket-binding motif, further supporting a role for RBP1 in bridging interactions between histone deacetylases and pRB/E2F complexes.

10 Example 8: RBP1 contains two independent transcriptional repression domains

As described herein, we identified a repression domain within RBP1. When Gal4 was fused to RBP1 having a deletion of this repression domain, referred to henceforth as R1, this mutant RBP1 product (Gal4-RBP1 $\Delta$ R1) still repressed CAT expression at high levels (Fig. 8A), suggesting that a second repression domain may exist towards the carboxyl terminus of RBP1. We therefore generated a series of in-frame carboxyl terminal deletion mutants that also lacked R1 (illustrated in Fig. 8B) and found that all, including Gal4- $\Delta$ R1-93C that lacked only 93 residues at the carboxyl terminus, failed to repress CAT expression (Fig. 8A). We showed that all mutants used in experiments in Fig. 8 were expressed at similar high levels. As these data indicated that the second repression domain likely lies at the very carboxyl terminus, we generated three additional constructs that contained only varying amounts of the carboxyl terminus of RBP1 linked to the Gal4DBD. Fig. 8A shows that all three, Gal4-R2(1311-C), Gal4-R2(1314-C) and Gal4-R2(1263-C), repressed CAT expression. Thus, a second RBP1 repression domain (termed R2) exists between residues 1167-1257. Furthermore, repression by neither R1 nor R2 relies on the LXCXE pocket-binding motif

-40-

domain (termed R2) exists between residues 1167-1257. Furthermore, repression by neither R1 nor R2 relies on the LXCXE pocket-binding motif when RBP1 is tethered to DNA by a heterologous DNA binding domain like Gal4DBD.

Example 9: Transcriptional repression by RBP1 is both dependent and  
5 independent of histone deacetylases activity

The ability of the pocket to actively repress transcription relies on both histone deacetylase-dependent and -independent mechanisms. Only a subset of promoters repressed by the pocket of pRB were sensitive to the specific histone deacetylase inhibitor, trichostatin A (TSA). Among these,  
10 the only Gal4-dependent promoter/reporter construct found to be repressed by the pocket and to be sensitive to TSA is G5MLPCAT (containing the adenovirus major late promoter). Fig. 8C shows that both Gal4-RBP1 and Gal4-pRB(pocket) repressed the G5MLPCAT reporter, as did both Gal4-R1 and Gal4-R2 that contain only R1 and R2, respectively. Fig. 8C also shows  
15 that following treatment of transfected cells with 330nM TSA for 24 hours prior to harvesting, repression by both Gal4-pRB(pocket) and Gal4-RBP1 was partially relieved; however, whereas drug treatment completely abolished repression by Gal4-R2, however, it had no effect on that by Gal4-R1. Thus it appears that repression by R2 depends on histone deacetylases whereas that by R1 does not.  
20 We also noted that whereas Gal4-HDAC1 repression activity is completely relieved by TSA (Fig. 8C), repression by Gal4-Ad5-E1B-55K, an adenoviral repressor known to block p53-dependent transactivation (Yew et al., Genes Dev. 8:190-202, 1994; Teodoro et al., J. Virol. 71:3620-3627, 1997), at this promoter is not affected by TSA, suggesting that the adenovirus major late promoter  
25 can also be subject to both histone deacetylase-dependent and -independent repression. These results therefore indicate that RBP1 plays an important role in repression by pRB as TSA only partially relieves repression by the pRB pocket. Interestingly, TSA had little effect on repression by

-41-

Gal4-RBP1 or Gal4-R2 using either the G5TKCAT or G5SV40CAT promoters. We extended studies from the synthetic G5MLPCAT construct to the E2F-dependent promoter regulating E2F-1 expression (E2F1-luc). Fig. 8D shows that both RBP1 and the pocket of pRB repressed expression of luciferase from the E2F1-luc reporter, whereas RBP1 lacking the LXCXE pocket-binding motif or a pRB point mutant (C706F) did not. Addition of TSA partially relieved repression by both RBP1 and the pRB pocket. These results indicate that both the RBP1 histone deacetylase-dependent and -independent repression activities can repress this E2F-dependent promoter via interactions with RB family members, and thus interactions with RBP1 could provide both types of repression activities attributed to the pocket of RB family members (Luo et al., Cell 92:463-473, 1998).

Example 10: Only one repression domain of RBP1 interacts with histone deacetylases

We tested the ability of individual R1 and R2 transcriptional repression domains to interact with histone deacetylases. Fig. 9A shows results of co-immunoprecipitation experiments using 293T cells expressing Gal4-RBP1 constructs and Flag-HDAC3 in which we immunoprecipitated RBP1 with anti-Gal4DBD antibodies, then detected HDAC3 by immunoblotting with anti-Flag antibody. We did not observe any HDAC3 binding using either R1 or just the ARID portion of R1, but such binding was clearly evident with both R2 constructs and with RBP1 lacking R1. Binding was eliminated or greatly reduced using RBP1 lacking R2. We obtained similar results using Flag-HDAC1 and -HDAC2.

At present it is not known if histone deacetylases bind directly to the RBP1 or if they interact indirectly via an additional R2-binding protein. It is highly likely that RBP1 directly binds to another member of the protein

-42-

complex that contains an HDAC. The most likely candidate is the SAP30 adaptor protein, but other proteins (e.g., SinA and SinB) that are part of protein complexes that contain HDACs may also bind directly to RBP1 (Pazin and Kadonaga, Cell 89:325-328, 1997, hereby incorporated by reference). The  
5 actual direct binding partners can be determined, for example, by ascertaining whether *in vitro* translated protein pairs are capable of direct binding. We have discovered the presence of several proteins that co-precipitate with RBP1 using anti-RBP1 antibodies. One of the most prominent species was a protein of about 48kDa. HDAC1 copurifies with RbAp48 that, along with RbAp46, plays  
10 a role in targeting HDAC1 to histones (Zhang et al., Mol Cell 1:1021-31, 1998). It is possible that both RBP1 and RbAp48/46 could co-exist in a single HDAC complex. RbAp48 binds to the so-called "extended pocket" including a carboxy terminal portion of pRB and thus might play a role as a linker for HDAC1. RbAp48 lacks the LXCXE- binding motif, however, and thus is not  
15 targeted to the small pocket. In addition, no interactions have been detected using the yeast two-hybrid system between the pRB pocket and HDAC1 or HDAC2, even though yeast cells contain high levels of MSI1, a protein that is highly homologous to RbAp48/46 (Zhang et al., Mol Cell 1:1021-31, 1998). It is very likely, then, that RBP1 is responsible for bridging the pocket of RB  
20 family members to histone deacetylase complexes to repress a diversity of E2F-dependent promoters. RBP1 therefore appears to represent a major component of the growth regulatory machinery controlled by RB family members.

The above-described results were obtained using the following materials  
25 and methods.

#### *Cell culture and transfection*

Monkey CV-1 cells (ATCC CCL 70) and human lung carcinoma-

-43-

H1299 cells were grown in Dulbecco's modified eagle medium (D-MEM) containing 10% fetal calf serum. Chinese hamster ovary (CHO) cells (ATCC CCL-61) were grown in  $\alpha$ -minimal essential medium supplemented with 10% fetal calf serum. Human cervical carcinoma (C33A; ATCC HTB-31), 293  
5 cells, 293T cells (a variant of 293 cells that express SV40 large T antigen) and Chinese hamster ovary (CHO) cells (ATCC CCL-61) were grown in  $\alpha$ -minimal essential medium supplemented with 10% fetal calf serum. In many experiments, CV-1 cells were induced to stop proliferating by incubating in D-MEM containing 2% dialyzed calf serum for 72 hrs, as described previously  
10 (Corbeil et al., *Oncogene* 15:657-668, 1997). Transfections were carried out by the calcium phosphate precipitation method (Graham and van der Eb, *Virology* 52:456-467, 1973) using pGEM plasmid as carrier DNA.

### *Plasmids*

E2F1-luc(wt) and E2F-luc (E2F-) reporter constructs were described  
15 previously (Smith et al., *Mol. Cell. Biol.* 16:6965-6976, 1996). Mammalian plasmid expression vectors included pCMVRBP1-HA (provided by Bill Kaelin) and pCMVRBP1 $\Delta$ LCE-HA (constructed by deleting residues 921-1056 of RBP1 coding region from pCMVRBP1-HA using restriction enzyme digests involving *NsiI* and *KpnI*). Plasmids expressing human pRB and p130, or the  
20 product of 12S Ad5 E1A mRNA were subcloned into pcDNA3 (Invitrogen). The G5TKCAT reporter construct has been described elsewhere (Yew et al., *Genes Dev.* 8:190-202, 1994; Teodoro et al., *J. Virol.* 71:3620-3627, 1997). Gal4-RBP1(SV40) plasmid was generated from pCMVRBP1-HA by polymerase chain reaction using Vent DNA polymerase (New England Biolabs,  
25 Mississauga, ON) with primers specific for the RBP1 coding region. The amplified products were digested with *SmaI* and *XbaI* restriction enzymes and subcloned in-frame with the Gal4 DNA binding domain of pSG424, which

-44-

expresses Gal4 fusion proteins in mammalian cells under the SV40 promoter. Gal4-RBP1(CMV) plasmid was constructed by subcloning the *HindIII* digested fragment of Gal4-RBP1(SV40) plasmid into pcDNA3, which expresses proteins in mammalian cells under the CMV promoter. All the carboxyl-terminal deletion mutants of Gal4-RBP1 were generated from restriction enzyme digests of the Gal4-RBP1(SV40) plasmid and subcloned into a modified pcDNA3 construct containing stop codons inserted 3' of the multicloning cassette. Gal4-RBP1-ΔLCE plasmid was constructed by deleting residues 921-1056 of RBP1 coding region from Gal4-RBP1(SV40) using restriction enzymes *NsiI* and *KpnI*. Gal4-RBP1-Rep plasmid was constructed by subcloning the restriction enzyme digested fragment that correspond to residues 388-599 of the Gal4-RBP1 coding region into pSG424. Gal4-Ad5E1B55K plasmid was described previously (Teodoro and Branton, J. Virol. 71:3620-3627, 1997). Mammalian expression plasmid expressing Gal4-pRB(pocket) was provided by Tony Kouzarides (Brehm et al., Nature 391:597-601, 1998). Flag-HDAC1, 2, and 3 and Gal4-HDAC1, HDAC2, HDAC3 constructs have been described elsewhere (Yang et al., Proc. Natl. Acad. Sci. USA 93:12845-12850, 1996; Yang et al., J. Biol. Chem. 272:28001-28007, 1997). Gal4-VP16 was provided by Arnie Berk (Yew et al., Nature 357:82-85, 1992). The G5MLPCAT reporter was provided by Doug Dean (Luo et al., Cell 92:463-473, 1998). Mutants of Gal4DBD-RBP1 were generated as follows: Gal4-ΔR1 mutants were generated using two specific primers close to the 3' end of the R1 region and the end of RBP1 coding sequence. PCR was done to generate fragments with unique restriction sites (*BspI* 107I, *HindIII*) which were subcloned into digested RBP1 constructs in which all RBP1 coding sequences between the beginning of R1 to the end of the protein had been removed using the same restriction enzymes. All carboxy terminal deletion mutants of Gal4-ΔR1 were generated from fragments

-45-

produced by restriction enzyme digestion of the Gal4- $\Delta$ R1 plasmid DNA and these were then subcloned into a modified pcDNA3 (Invitrogen) construct containing stop codons inserted 3' of the multicloning cassette. Gal4-R2 truncation mutants were constructed by subcloning the restriction  
5 enzyme-digested fragments that correspond to residues 1311-1404, or 1314-1404, or 1263-1404 of the Gal4-RBP1 coding region into pSG424. Mammalian expression plasmids encoding the pocket of pRB and the inactive pRB pocket (C706F) mutant have been described previously (Kaelin et al., Cell 70:351-364, 1992). All constructs made were confirmed by sequencing using  
10 specific primers.

*Cell Lysis and electrophoretic mobility shift assay(EMSA)*

CV-1 cells were harvested and whole cell extracts were prepared using lysis buffer (50 mM HEPES-KOH, pH 7.9, containing 0.4 M KCl, 0.1%(v/v) NP-40, 4 mM NaF, 4 mM NaVO<sub>4</sub>, 0.2 mM EDTA, 0.2 mM EGTA, 10%  
15 Glycerol (v/v), 1 mM DTT, 0.5 mM PMSF, 1  $\mu$ g/mL pepstatin, 1  $\mu$ g/mL leupeptin and 1  $\mu$ g/mL aprotinin) and centrifuged for 45 min at 20,000 x g (Corbeil et al., Oncogene 11:909-920, 1995; Corbeil et al., Oncogene 15:657-668, 1997). EMSA was performed using whole cell extracts and an E2F-specific oligonucleotide (GATTTAAGTTTCGCGCCCTTTCTCAA (SEQ  
20 ID NO: 1)) and double stranded oligonucleotides were labeled using [ $\alpha^{32}$ P]dCTP and Klenow polymerase. EMSA was carried out in 20  $\mu$ L reaction mixtures containing 10-20  $\mu$ g of whole cell extract in 20 mM HEPES-KOH (pH 7.9) containing 40 mM KCl, 1 mM MgCl<sub>2</sub>, 0.1 mM EGTA, 0.4 mM DTT, 4  $\mu$ g BSA, 2.5% (v/v) Ficoll, 1-2  $\mu$ g of salmon sperm DNA, and labeled probe  
25 containing 20,000-50,000 cpm. Antibody supershift experiments were performed by incubating extracts for 10 min on ice with 1-2  $\mu$ g of antisera recognizing p130, pRB, p107, or RBP1. Reactions were incubated at room



-46-

temperature for 30 min and then samples were loaded on a 4% polyacrylamide gel (30:1 polyacrylamide: bis-acrylamide) in 0.25X TBE and subjected to electrophoresis at 200 V for 2 hr. Gels were dried and analyzed by autoradiography using Kodak X-omat film.

## 5 *Antibodies*

A rabbit polyclonal antibody raised against HDAC2 was described previously (Laherty et al., Cell 89:349-356, 1997). Antibodies against HDAC1 and HDAC3 were prepared against peptides corresponding to the unique carboxy termini of HDAC1 (EEKPEAKGVKEEVKLA) (SEQ ID NO: 2) and  
10 HDAC3 (NEFYDGDHDNDKESDVEI) (SEQ ID NO: 3) coupled to KLH and injected separately into New Zealand white rabbits. The resulting antibodies were immunoaffinity purified on peptide columns. Anti-pRB antibody G3-245 was purchased from PharMingen (San Diego, CA). Antibodies against p130 (C-20), p107 (C-18), and Gal4DBD (RK5C1) were purchased from Santa Cruz,  
15 anti-HA antibody HA.11 was purchased from BAbCo (Richmond, CA) and anti-Flag antibody M2 from Sigma, (St. Louis, MO).

## *Chloramphenicol acetyltransferase (CAT), $\beta$ -galactosidase and luciferase assays*

CAT and  $\beta$ -galactosidase assays were performed as described previously  
20 (Teodoro and Branton, J. Virol. 71:3620-3627, 1997). Luciferase assays were performed as described elsewhere (Wrana et al., Cell 71:1003-1014, 1992).

## *Colony formation assay*

CV-1 cells were transfected with 3  $\mu$ g of DNA from plasmids expressing neo as well as RBP1, including pRcCMVRBP1-HA (encoding an  
25 HA-tagged version of RBP1), pcDNA3Gal4-RBP1, and pcDNA3Gal4-RBP1 $\Delta$ LCE. Cells were plated in medium containing 500  $\mu$ g/mL of G418,

-47-

allowed to grow for two weeks, and then stained with crystal violet.

#### *Binding assays*

- 1  $\mu$ g of cDNA encoding Gal4DBD, fused with the pocket of pRB, RBP1, or mutated RBP1, were introduced using lipofectamine (NEN Life
- 5 Science, Boston, MA) along with 1  $\mu$ g of those encoding Flag-tagged HDAC1, HDAC2, or HDAC3 into H1299 or 293T cells. In some experiments, 1  $\mu$ g of cDNAs encoding HA- tagged RBP1 or RBP1 mutants were introduced using lipofectamine along with 1  $\mu$ g of those encoding Gal4DBD fused HDAC1, HDAC2, or HDAC3 into H1299 or 293T. Cells were harvested 40 hours
- 10 post-transfection and lysed with "low stringency buffer." Cell extracts were diluted to 150mM KCl in a 1 mL volume and precleared with protein G-Sepharose (Pharmacia, Baie D'Urfe, QC) for 2 hours. Precleared extracts were incubated with 1  $\mu$ g of RK5C1-Gal4DBD antibody (Santa Cruz) and 30  $\mu$ L of a 50% slurry of protein G-Sepharose for at least 12 hours.
- 15 Immunoprecipitated Gal4-tagged protein complexes were washed six times with lysis buffer and eluted by boiling in 2X sample buffer. Eluted proteins were subjected to SDS-PAGE using a 10% polyacrylamide gel, and proteins were then transferred to PVDF membranes (Millipore, Bedford, MA) which were probed with either anti-HA or anti-Flag antibody and then by
- 20 HRP-conjugated goat anti-mouse (light chain specific) secondary antibody. Binding was detected by Enhanced Luminol Reagent (NEN Life Science).

#### Other Embodiments

- All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the
- 25 described invention will be apparent to those skilled in the art. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not

-48-

be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the invention.

5           What is claimed is:

-49-

1. A method for modulating proliferation of a cell, said method comprising contacting said cell with an RBP1 polypeptide or a compound that modulates RBP1 biological activity or expression.
2. The method of claim 1, wherein said cell proliferation is decreased.
- 5       3. A method for identifying a compound that modulates cell proliferation, said method comprising contacting a cell with a candidate compound and monitoring the level of expression of an RBP1 polypeptide in said cell, wherein a change in said level of expression in response to said candidate compound, relative to the level of expression in a cell not contacted  
10       with said candidate compound, identifies said compound as one that modulates cell proliferation.
4. A method for identifying a compound that modulates cell proliferation, said method comprising:
  - (a) providing an RBP1 polypeptide having biological activity;
  - 15       (b) contacting said RBP1 polypeptide with a candidate compound; and
  - (c) monitoring the level of said biological activity of said RBP1 polypeptide, wherein a change in said level of biological activity in response to said candidate compound, relative to the level of biological activity of an RBP1 polypeptide not contacted with said candidate compound, identifies said  
20       compound as a compound that modulates cell proliferation.
5. The method of claim 4, wherein said RBP1 polypeptide is in a cell.
6. The method of claim 1 or 5, wherein said cell is in a mammal.

-50-

7. The method of claim 6, wherein said mammal is a human or a rodent.

8. The method of claim 4, wherein said RBP1 polypeptide is in a cell-free system.

9. A method for determining whether a patient has a cell proliferation  
5 disease or an increased likelihood of developing said disease, said method  
comprising measuring the level of RBP1 biological activity in a sample from  
the mammal, wherein an alteration in said level of said biological activity,  
relative to a level of RBP1 biological activity in a matched sample from an  
unaffected person, indicates that the patient has said disease or increased  
10 likelihood of developing said disease.

10. The method of claim 1, 4, or 9, wherein said biological activity is  
binding of said RBP1 polypeptide to an E2F protein or repression of E2F-  
mediated transcription.

11. The method of claim 10, wherein said E2F is E2F-1, E2F-2, E2F-3,  
15 E2F-4, or E2F-5.

12. The method of claim 1, 4, or 9, wherein said biological activity is  
binding of said RBP1 polypeptide to an HDAC or a protein that is part of a  
complex containing an HDAC.

13. The method of claim 12, wherein said protein SAP30, SinA, or  
20 SinB.

14. A method for determining whether a patient has a cell proliferation

-51-

disease or an increased likelihood of developing said disease, said method comprising measuring the level of expression of an RBP1 polypeptide in a sample from said mammal, wherein an alteration in said level of expression of said RBP1 polypeptide, relative to a matched sample from an unaffected  
5 person, indicates that the patient has a cell proliferation disease or increased likelihood of developing said disease.

15. A method of repressing transcription in a cell, said method comprising introducing into said cell a polypeptide comprising: (i) a repressor domain from RBP1, and (ii) a DNA binding domain.

10 16. A polypeptide comprising: (i) a repressor domain from RBP1, and (ii) a DNA binding domain.

17. The polypeptide of claim 16, wherein said repressor domain comprises amino acids 241-452 or 1167-1257 of SEQ ID NO: 4.

15 18. A nucleic acid molecule encoding a polypeptide comprising: (i) a repressor domain from RBP, and (ii) a DNA binding domain.

19. A composition comprising (i) a substantially purified RBP1 polypeptide, a substantially purified nucleic acid encoding an RBP1 polypeptide, or a substantially purified antibody that specifically binds an RBP1 polypeptide, and (ii) a physiologically acceptable carrier.

20 20. A kit for determining whether a human has a cell proliferation disease or an increased likelihood of developing said disease, said kit comprising a substantially pure antibody that specifically binds an RBP1

polypeptide.

21. The kit of claim 20, wherein said kit further comprises a means for detecting said binding of said antibody to said RBP1 polypeptide.

22. Use of an RBP1 polypeptide, a nucleic acid molecule encoding an  
5 RBP1 polypeptide, or an antibody that specifically binds to an RBP1  
polypeptide for preparing a medicament for the treatment of a cell proliferation  
disease.






Fig. 1




Fig. 2A




Fig. 2B




Fig. 2C




Fig. 3A

4/20

pcDNA3




Fig. 3B

pRcCMV-RBP1-HA



Fig. 3C

5/20

pcDNA3Gal4-RBP1-dl-747 C




Fig. 3D

pcDNA3Gal4-RBP1




Fig. 3E

6/20




Fig. 4A

7/20




Fig. 4B



8/20




Fig. 5A





Fig. 5B


9/20

**H1299 cells (p53<sup>-/-</sup>)****Co-transfection of Gal4Rb(pocket)/Flag-HDAC**

Gal4-Rb	-	-	-	+	+	+
Gal4-VP16	+	+	+	-	-	-
Flag-HDAC1	+	-	-	+	-	-
Flag-HDAC2	-	+	-	-	+	-
Flag-HDAC3	-	-	+	-	-	+

**Fig. 6A****293T cells (E1A & Tag)****Co-transfection of Gal4Rb(pocket)/Flag-HDAC**


Gal4-Rb	+	+	+	+	+	+
Flag-HDAC1	+	+	-	-	-	-
Flag-HDAC2	-	-	+	+	-	-
Flag-HDAC3	-	-	-	-	+	+

**Fig. 6B**

10/20

**293 cells (E1A)**Co-transfection of **Gal4Rb(pocket)** or **Gal4RBP1/Flag-HDAC**

Gal4-RBP1	+	-	+	-	+	-
Gal4-Rb	-	+	-	+	-	+
Flag-HDAC1	+	+	-	-	-	-
Flag-HDAC2	-	-	+	+	-	-
Flag-HDAC3	-	-	-	-	+	+



**Fig. 6C**

11/20


**H1299 cells preincubated with GST-E1A**

Co-transfection of Gal4Rb(pocket) /Flag-HDAC1

Gal4	+	-	-	-	-	-	-
Gal4-VP16	-	+	-	-	-	-	-
Flag-HDAC1	+	+	+	+	+	+	+
Gal4-Rb	-	-	+	+	+	+	+
GST	-	-	+	-	-	-	-
GST-E1A	-	-	-	-	-	-	▲





GST	-	-	+
GST-E1A	-	+	-
Glutathione sepharose	+	+	+

**Fig. 6D**


12/20

**H1299 cells (p53<sup>-/-</sup>)****Co-transfection of Gal4RBP1/Flag-HDAC**

Gal4-RBP1	-	-	-	+	+	+
Gal4-VP16	+	+	+	-	-	-
Flag-HDAC1	+	-	-	+	-	-
Flag-HDAC2	-	+	-	-	+	-
Flag-HDAC3	-	-	+	-	-	+

**Fig. 7A****293T cells (E1A & Tag)****Co-transfection of Gal4RBP1/Flag-HDAC**


Gal4-RBP1	+	+	+	+	+	+
Flag-HDAC1	+	+	-	-	-	-
Flag-HDAC2	-	-	+	+	-	-
Flag-HDAC3	-	-	-	-	+	+

**Fig. 7B**


13/20

**293T cells (E1A & Tag)****Co-transfection of RBP1-HA/Gal4-HDAC**

Gal4	+	-	-	-	-
Gal4-VP16	-	+	-	-	-
Gal4-HDAC1	-	-	+	-	-
Gal4-HDAC2	-	-	-	+	-
Gal4-HDAC3	-	-	-	-	+
RBP1-HA	+	+	+	+	+

**Fig. 7C**

14/20








15/20

**293T cells (E1A & Tag)****Co-transfection of RBP1dl-LXCXE-HA/Gal4-HDAC**

RBP1dl-LXCXE-HA	+	+	+	+
Gal4-VP16	+	-	-	-
Gal4-Rb	-	+	-	-
Gal4-HDAC1	-	-	+	-
Gal4-HDAC2	-	-	-	+

**Co-transfection of Gal4RBP1dl-LXCXE/Flag-HDAC**

Gal4-VP16	+	-	-	-	+	-	-	-
Gal4-Rb	-	+	-	-	-	+	-	-
Gal4-RBP1	-	-	+	-	-	-	+	-
Gal4-RBP1dl-LXCXE	-	-	-	+	-	-	-	+
Flag-HDAC1	+	+	+	+	-	-	-	-
Flag-HDAC3	-	-	-	-	+	+	+	+

**Fig. 7E**

16/20




Fig. 8A

17/20




Fig. 8B




Fig. 8C




Fig. 8D

20/20

293T cells (E1A &amp; Tag)


Co-transfection of **Gal4RBP1** mutants/Flag-HDAC3

Fig. 9

## SEQUENCE LISTING

&lt;110&gt; GeminX, Inc.

&lt;120&gt; RBP1 POLYPEPTIDES AND USES THEREOF

&lt;130&gt; 50013/018WO2

&lt;150&gt; 60/143,315

&lt;151&gt; 1999-07-12

&lt;160&gt; 4

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthetic based on Homo sapiens

&lt;400&gt; 1

gatttaagtt tcgcgccctt tctcaa

26

&lt;210&gt; 2

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthetic based on Homo sapiens

&lt;400&gt; 2

Glu	Glu	Lys	Pro	Glu	Ala	Lys	Gly	Val	Lys	Glu	Glu	Val	Lys	Leu	Ala
1				5				10					15		

&lt;210&gt; 3

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthetic based on Homo sapiens

&lt;400&gt; 3

Asn	Glu	Phe	Tyr	Asp	Gly	Asp	His	Asp	Asn	Asp	Lys	Glu	Ser	Asp	Val
1				5				10						15	

Glu Ile

&lt;210&gt; 4

&lt;211&gt; 1257

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

Met	Lys	Ala	Ala	Asp	Glu	Pro	Ala	Tyr	Leu	Thr	Val	Gly	Thr	Asp	Val
1				5					10					15	
Ser	Ala	Lys	Tyr	Arg	Gly	Ala	Phe	Cys	Glu	Ala	Lys	Ile	Lys	Thr	Val
			20					25						30	
Lys	Arg	Leu	Val	Lys	Val	Lys	Val	Leu	Leu	Lys	Gln	Asp	Asn	Thr	Thr
		35					40						45		
Gln	Leu	Val	Gln	Asp	Asp	Gln	Val	Lys	Gly	Pro	Leu	Arg	Val	Gly	Ala
	50					55					60				
Ile	Val	Glu	Thr	Arg	Thr	Ser	Asp	Gly	Ser	Phe	Gln	Glu	Ala	Ile	Ile
65					70					75					80
Ser	Lys	Leu	Thr	Asp	Ala	Ser	Trp	Tyr	Thr	Val	Val	Phe	Asp	Asp	Gly
				85					90					95	
Asp	Glu	Arg	Thr	Leu	Arg	Arg	Thr	Ser	Leu	Cys	Leu	Lys	Gly	Glu	Arg
			100					105						110	
His	Phe	Ala	Glu	Ser	Glu	Thr	Leu	Asp	Gln	Leu	Pro	Leu	Thr	Asn	Pro
		115					120							125	
Glu	His	Phe	Gly	Thr	Pro	Val	Ile	Ala	Lys	Lys	Thr	Asn	Arg	Gly	Arg
	130					135								140	
Arg	Ser	Ser	Leu	Pro	Val	Thr	Glu	Asp	Glu	Lys	Glu	Glu	Glu	Ser	Ser
145					150					155					160
Glu	Glu	Glu	Asp	Glu	Asp	Lys	Arg	Arg	Leu	Asn	Asp	Glu	Leu	Leu	Gly
				165					170					175	
Lys	Val	Val	Ser	Val	Val	Ser	Ala	Thr	Glu	Arg	Thr	Glu	Trp	Tyr	Pro
			180					185						190	
Ala	Leu	Val	Ile	Ser	Pro	Ser	Cys	Asn	Asp	Asp	Ile	Thr	Val	Lys	Lys
		195					200							205	
Asp	Gln	Cys	Leu	Val	Arg	Ser	Phe	Ile	Asp	Ser	Lys	Phe	Tyr	Ser	Ile
	210				215						220				
Ala	Arg	Lys	Asp	Ile	Lys	Glu	Val	Asp	Ile	Leu	Asn	Leu	Pro	Glu	Ser
225					230					235					240
Glu	Leu	Ser	Thr	Lys	Pro	Gly	Leu	Gln	Lys	Ala	Ser	Ile	Phe	Leu	Lys
				245					250					255	
Thr	Arg	Val	Val	Pro	Asp	Asn	Trp	Lys	Met	Asp	Ile	Ser	Glu	Ile	Leu
		260						265						270	
Glu	Ser	Ser	Ser	Asp	Asp	Glu	Asp	Gly	Pro	Ala	Glu	Glu	Asn	Asp	
		275				280					285				
Glu	Glu	Lys	Glu	Lys	Glu	Ala	Lys	Lys	Thr	Glu	Glu	Glu	Val	Pro	Glu
	290					295					300				
Glu	Glu	Leu	Asp	Pro	Glu	Glu	Arg	Asp	Asn	Phe	Leu	Gln	Gln	Leu	Tyr
305					310					315					320
Lys	Phe	Met	Glu	Asp	Arg	Gly	Thr	Pro	Ile	Asn	Lys	Pro	Pro	Val	Leu
				325					330					335	
Gly	Tyr	Lys	Asp	Leu	Asn	Leu	Phe	Lys	Leu	Phe	Arg	Leu	Val	Tyr	His
			340					345					350		
Gln	Gly	Gly	Cys	Asp	Asn	Ile	Asp	Ser	Gly	Ala	Val	Trp	Lys	Gln	Ile
	355					360						365			
Tyr	Met	Asp	Leu	Gly	Ile	Pro	Ile	Leu	Asn	Ser	Ala	Ala	Ser	Tyr	Asn
	370					375					380				
Leu	Lys	Thr	Ala	Tyr	Arg	Lys	Tyr	Leu	Tyr	Gly	Phe	Glu	Glu	Tyr	Cys
385					390					395					400



Arg Ser Ala Asn Ile Gln Phe Arg Thr Val His His His Glu Pro Lys  
 405 410 415  
 Val Lys Glu Glu Lys Lys Asp Leu Glu Glu Ser Met Glu Glu Ala Leu  
 420 425 430  
 Lys Leu Asp Gln Glu Met Pro Leu Thr Glu Val Lys Ser Glu Pro Glu  
 435 440 445  
 Glu Asn Ile Asp Ser Asn Ser Glu Ser Glu Arg Glu Glu Ile Glu Leu  
 450 455 460  
 Lys Ser Pro Arg Gly Arg Arg Arg Ile Ala Arg Asp Val Asn Ser Ile  
 465 470 475 480  
 Lys Lys Glu Ile Glu Glu Glu Lys Thr Glu Asp Lys Leu Lys Asp Asn  
 485 490 495  
 Asp Thr Glu Asn Lys Asp Val Asp Asp Asp Tyr Glu Thr Ala Glu Lys  
 500 505 510  
 Lys Glu Asn Glu Leu Leu Leu Gly Arg Lys Asn Thr Pro Lys Gln Lys  
 515 520 525  
 Glu Lys Lys Ile Lys Lys Gln Glu Asp Ser Asp Lys Asp Ser Asp Glu  
 530 535 540  
 Glu Glu Glu Lys Ser Gln Glu Arg Glu Glu Thr Glu Ser Lys Cys Asp  
 545 550 555 560  
 Ser Glu Gly Glu Glu Asp Glu Glu Asp Met Glu Pro Cys Leu Thr Gly  
 565 570 575  
 Thr Lys Val Lys Val Lys Tyr Gly Arg Gly Lys Thr Gln Lys Ile Tyr  
 580 585 590  
 Glu Ala Ser Ile Lys Ser Thr Glu Ile Asp Asp Gly Glu Val Leu Tyr  
 595 600 605  
 Leu Val His Tyr Tyr Gly Trp Asn Val Ser Tyr Asp Glu Trp Val Lys  
 610 615 620  
 Ala Asp Arg Ile Ile Trp Pro Leu Asp Lys Gly Gly Pro Lys Lys Lys  
 625 630 635 640  
 Gln Lys Lys Lys Ala Lys Asn Lys Glu Asp Ser Glu Lys Asp Glu Lys  
 645 650 655  
 Arg Asp Glu Glu Arg Gln Lys Ser Lys Arg Gly Arg Pro Pro Leu Lys  
 660 665 670  
 Ser Thr Leu Ser Ser Asn Met Pro Tyr Gly Leu Ser Lys Thr Ala Asn  
 675 680 685  
 Ser Glu Gly Lys Ser Asp Ser Cys Ser Ser Asp Ser Glu Thr Glu Asp  
 690 695 700  
 Ala Leu Glu Lys Asn Leu Ile Asn Glu Glu Leu Ser Leu Lys Asp Glu  
 705 710 715 720  
 Leu Glu Lys Asn Glu Asn Leu Asn Asp Asp Lys Leu Asp Glu Glu Asn  
 725 730 735  
 Pro Lys Ile Ser Ala His Ile Leu Lys Glu Asn Asp Arg Thr Gln Met  
 740 745 750  
 Gln Pro Leu Glu Thr Leu Lys Leu Glu Val Gly Glu Asn Glu Gln Ile  
 755 760 765  
 Val Gln Ile Phe Gly Asn Lys Met Glu Lys Ala Glu Glu Val Lys Lys  
 770 775 780  
 Glu Ala Glu Lys Ser Pro Lys Gly Lys Gly Arg Arg Ser Lys Thr Lys  
 785 790 795 800  
 Asp Leu Ser Leu Glu Ile Ile Lys Ile Ser Ser Phe Gly Gln Asn Glu  
 805 810 815  
 Ala Gly Ser Glu Pro His Ile Glu Ala His Ser Leu Glu Leu Ser Ser  
 820 825 830  
 Leu Asp Asn Lys Asn Phe Ser Ser Ala Thr Glu Asp Glu Ile Asp Gln

835	840	845
Cys Val Lys Glu Lys Lys Leu Lys Arg Lys Ile Leu Gly Gln Ser Ser		
850	855	860
Pro Glu Lys Lys Ile Arg Ile Glu Asn Gly Met Glu Met Thr Asn Thr		
865	870	875
Val Ser Gln Glu Arg Thr Ser Asp Cys Ile Gly Ser Glu Gly Met Lys		880
	885	890
Asn Leu Asn Phe Glu Gln His Phe Glu Arg Glu Asn Glu Gly Met Pro		895
	900	905
Ser Leu Ile Ala Glu Ser Asn Gln Cys Ile Gln Gln Leu Thr Ser Glu		910
	915	920
Arg Phe Asp Ser Pro Ala Glu Glu Thr Val Asn Ile Pro Leu Lys Glu		925
	930	935
Asp Glu Asp Ala Met Pro Leu Ile Gly Pro Glu Thr Leu Val Cys His		940
945	950	955
Glu Val Asp Leu Asp Asp Leu Asp Glu Lys Asp Lys Thr Ser Ile Glu		960
	965	970
Asp Val Ala Val Glu Ser Ser Glu Ser Asn Ser Leu Val Ser Ile Pro		975
	980	985
Pro Ala Leu Pro Pro Val Val Gln His Asn Phe Ser Val Ala Ser Pro		990
	995	1000
Leu Thr Leu Ser Gln Asp Glu Ser Arg Ser Val Lys Ser Glu Ser Asp		1005
1010	1015	1020
Ile Thr Ile Glu Val Asp Ser Ile Ala Glu Glu Ser Gln Glu Gly Leu		
1025	1030	1035
Cys Glu Arg Glu Ser Ala Asn Gly Phe Glu Thr Asn Val Ala Ser Gly		104
	1045	1050
Thr Cys Ser Ile Ile Val Gln Glu Arg Glu Ser Arg Glu Lys Gly Gln		1055
	1060	1065
Lys Arg Pro Ser Asp Gly Asn Ser Gly Leu Met Ala Lys Lys Gln Lys		1070
	1075	1080
Arg Thr Pro Lys Arg Thr Ser Ala Ala Ala Lys Asn Glu Lys Asn Gly		1085
1090	1095	1100
Thr Gly Gln Ser Ser Asp Ser Glu Asp Leu Pro Val Leu Asp Asn Ser		
1105	1110	1115
Ser Lys Cys Thr Pro Val Lys His Leu Asn Val Ser Lys Pro Gln Lys		112
	1125	1130
Leu Ala Arg Ser Pro Ala Arg Ile Ser Pro His Ile Lys Asp Gly Glu		1135
	1140	1145
Lys Asp Lys His Arg Glu Lys His Pro Asn Ser Ser Pro Arg Thr Tyr		1150
	1155	1160
Lys Trp Ser Phe Gln Leu Asn Glu Leu Asp Asn Met Asn Ser Thr Glu		1165
1170	1175	1180
Arg Ile Ser Phe Leu Gln Glu Lys Leu Gln Glu Ile Arg Lys Tyr Tyr		
1185	1190	1195
Met Ser Leu Lys Ser Glu Val Ala Thr Ile Asp Arg Arg Arg Lys Arg		120
	1205	1210
Leu Lys Lys Lys Asp Arg Glu Val Ser His Ala Gly Ala Ser Met Ser		1215
	1220	1225
Ser Ala Ser Ser Asp Thr Gly Met Ser Pro Ser Ser Ser Ser Pro Pro		1230
	1235	1240
Gln Asn Val Leu Ala Val Glu Cys Arg		1245
1250	1255	

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 99/01690

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C07K16/18 A61K38/17 A61K39/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ALI R FATTAEY ET AL: "Characterization of the retinoblastoma binding proteins RBP1 and RBP2" ONCOGENE, GB, BASINGSTOKE, HANTS, vol. 8, no. 11, November 1993 (1993-11), pages 3149-3156-3156, XP002107565 ISSN: 0950-9232 cited in the application the whole document	1-22
X	DEFEO-JONES, DEBORAH ET AL: "Cloning of cDNAs for cellular proteins that bind to the retinoblastoma gene product" NATURE (LONDON), 1991, 352, 251-4, XP000872842 cited in the application the whole document	1-22
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"S" document member of the same patent family

Date of the actual completion of the international search

11 February 2000

Date of mailing of the international search report

24/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5518 Patentplan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hillenbrand, G

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 99/01690

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>OTTERSON GA ET AL: "Alternative splicing of the *RBP1* gene clusters in an internal exon that encodes potential phosphorylation sites." ONCOGENE, APR 1993, 8 (4) P949-57, XP000872340 ENGLAND cited in the application the whole document</p>	1-22
X	<p>LAI A ET AL: "*RBP1* induces growth arrest by repression of E2F-dependent transcription." ONCOGENE, MAR 25 1999, 18 (12) P2091-100, XP000872491 ENGLAND the whole document</p>	1-22
A	<p>WO 93 15227 A (UNIV DUKE) 5 August 1993 (1993-08-05) the whole document</p>	1-22

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 99/01690

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9315227 A	05-08-1993	AU 3609693 A US 5650287 A	01-09-1993 22-07-1997